

Structure functions and energy dissipation dependence on Reynolds number

G. Boffetta

Dipartimento di Fisica Generale and INFM, Università di Torino, Via Pietro Giuria 1, I-10125 Torino, Italy

G. P. Romano

Dipartimento di Meccanica ed Aeronautica, Università di Roma "la Sapienza," Via Eudossiana 18, I-00184 Roma, Italy

(Received 13 July 2001; accepted 10 July 2002; published 3 September 2002)

The dependence of the statistics of energy dissipation on the Reynolds number is investigated in an experimental jet flow. In a range of about one decade of Re_λ (from about 200 to 2000) the adimensional mean energy dissipation is found to be independent on Re_λ , while the higher moments of dissipation show a power-law dependence. The scaling exponents are found to be consistent with a simple prediction based on the multifractal model for inertial range structure functions. This is an experimental confirmation of the connection between inertial range quantities and dissipation statistics predicted by the multifractal approach. © 2002 American Institute of Physics.

[DOI: 10.1063/1.1504449]

I. INTRODUCTION

There is considerable experimental¹ and numerical² evidence that the longitudinal velocity difference structure functions

$$S_q(\ell) \equiv \langle \delta u(\ell)^q \rangle = C_q \langle \epsilon \rangle^{q/3} \ell^{q/3} \left(\frac{\ell}{L} \right)^{\zeta_q - q/3}, \quad (1)$$

are affected by intermittency corrections in the scaling exponents ζ_q which deviate from the Kolmogorov self-similar prediction $\zeta_q = q/3$.^{3,4} (C_q is a constant possibly depending on the Reynolds number, ϵ is the turbulent kinetic energy dissipation, ℓ is a generic length scale and L is the integral length scale). The estimate of ζ_q is now available for a wide range of Reynolds numbers and different flow configurations. The analysis of experimental data shows that structure functions display the scaling behavior (1) for sufficiently high Reynolds numbers, Re , and that in such conditions the longitudinal exponents ζ_p become independent on Re .^{5,6} There is also experimental support for the scaling exponents to be Re -independent also at small Re [when the scaling behavior in (1) is even not observable] if one make use of the so-called extended self-similarity analysis (ESS).⁷ However, the precise relation between this empirical result and the classical scaling behavior (1), expected from dimensional arguments, is not well understood. We will not further discuss this point here.

The basic fundamental property of fully developed turbulence is that the average energy dissipation $\langle \epsilon \rangle$ entering in (1) is asymptotically Re -independent, when adimensionalized with large scale variables.^{3,8,9} On the other hand, it is well known that the statistics of the local energy dissipation, which assuming isotropy can be defined in terms of its one-dimensional (1D) surrogate

$$\epsilon(x) = 15\nu \left(\frac{\partial u}{\partial x} \right)^2, \quad (2)$$

strongly depends on Re , becoming more and more intermittent with increasing Re .^{10,11} This effect is also reflected in the tails of the pdf of velocity increments at very small separations.¹²⁻¹⁴ A statistical description of the behavior of the local energy dissipation (2) can be obtained in terms of quantities defined in the inertial range, such as (1), within the multifractal approach. This approach, originally introduced as a phenomenological model for the inertial range statistics,^{15,16} has been extended to the prediction of dissipative scale statistics.^{4,10,11,17,18} In particular, the multifractal model predicts a scaling behavior of the moments of (2) with Re ,¹⁹ which has been recently measured in a simplified model of turbulence.²⁰

In the present paper we investigate the Reynolds dependence of the statistics of (2) in a experimental water jet. We have examined a series of data obtained from laser Doppler anemometer (LDA) measurements which cover about one decade of Reynolds numbers. The quality of the statistics allows us to compute high-order structure functions with reasonable accuracy and to partially resolve the dissipative scales. Our main result is that the dependence of the statistics of dissipation on the Reynolds number is found to be consistent with the multifractal prediction obtained by assuming a fluctuating dissipative scale.

The remaining of the paper is organized as follows. In Sec. II we introduce the theoretical models for the statistics of dissipation. In Sec. III, the experimental setup and the data analysis procedure. Section IV is devoted to the presentation of the results, whereas concluding remarks are given in Sec. V.

II. MULTIFRACTAL DESCRIPTION OF ENERGY DISSIPATION

A phenomenological description of intermittency is given by the multifractal model.¹⁵ This model introduces a continuous set of scaling exponents h which locally relates the velocity fluctuations at scale ℓ entering in (1) with a large-scale velocity fluctuation u'

$$\delta u(\ell) \sim u' \left(\frac{\ell}{L} \right)^h \tag{3}$$

The local exponent h is realized with a probability which scales with $(\ell/L)^{Z(h)}$ where $Z(h)$ is the codimension of the fractal set on which the h -scaling holds. The scaling exponent of structure functions (1) are obtained by a steepest descent argument over h

$$\zeta_q = \inf_h [qh + Z(h)]. \tag{4}$$

The scaling region of (3) is bounded from below by the Kolmogorov dissipative scale η at which dissipation starts to dominate, i.e., at which the local Reynolds number is of order 1

$$\frac{\eta \delta u(\eta)}{\nu} \simeq 1. \tag{5}$$

From (3) and (5) one obtains that in the multifractal description of intermittency the dissipative scale is a fluctuating quantity, i.e., depends on the local scaling exponent h according to

$$\eta \sim L \left(\frac{u' L}{\nu} \right)^{-1/(1+h)} \sim L \text{Re}_\lambda^{-2/(1+h)}, \tag{6}$$

where $\text{Re}_\lambda = u' \lambda / \nu$ is the Reynolds number based on the Taylor microscale $\lambda = \sqrt{15 \nu u'^2 / \langle \epsilon \rangle}$.

Below the dissipative scale the flow can be assumed smooth and one can replace the derivative in (2) with

$$\epsilon = 15 \nu \left(\frac{\delta u(\eta)}{\eta} \right)^2. \tag{7}$$

Assuming that the multifractal model can be pushed down to the dissipative scale, one can evaluate the statistics of (2) by inserting (3) and (6) into (7). One ends with the expression^{4,19,20}

$$\begin{aligned} \langle \epsilon^p \rangle &\sim \langle \epsilon \rangle^p \int d\mu(h) \text{Re}_\lambda^{-2[3ph-p+Z(h)]/(1+h)} \\ &\sim \langle \epsilon \rangle^p \text{Re}_\lambda^{-2\theta_p}, \end{aligned} \tag{8}$$

where again the integral has been evaluated by a steepest descent argument as

$$\theta_p = \inf_h \left[\frac{3ph - p + Z(h)}{1+h} \right]. \tag{9}$$

The standard inequality in the multifractal model (following from the exact result $\zeta_3 = 1$), $Z(h) \geq 1 - 3h$,⁴ implies for (9) $\theta_1 = 0$ which is nothing but the request of finite non-

vanishing dissipation in the limit $\text{Re} \rightarrow \infty$. For $p > 1$, $\theta_p < 0$, i.e., the tail of the distribution of ϵ becomes wider with the Reynolds number.

Let us remark that the prediction (9) is based on the assumption of a fluctuating dissipative scale η according to (6). If one assume, on the contrary, that dissipation enters in (7) as an average quantity one ends up with a different predictions for the exponents θ_p ²¹ [see below Eq. (13)]. Numerical simulations with a simplified model of turbulent cascade have shown that the exponents (6) are indeed observed and the alternative prediction is ruled out.²⁰ In the next section, we will see that also our experimental data are in agreement with prediction (9).

In the following we will consider data analysis of one component of the velocity in water jets at different Reynolds numbers. Because of the discretization of the acquisition, we are forced to replace spatial derivatives with velocity differences at small scales. The key quantity for our discussion is a generalization of (2) over a finite scale ℓ

$$E(\ell) \equiv 15 \nu \left(\frac{\delta u(\ell)}{\ell} \right)^2. \tag{10}$$

$E(\ell)$ is a convenient definition of surrogate energy dissipation if the scale ℓ is sufficiently small. The average dissipative scale dependence on Reynolds number can be obtained from (7) and (1) as

$$\bar{\eta} \simeq L \text{Re}_\lambda^{-2/(2-\zeta_2)}. \tag{11}$$

In the analysis of experimental data it is convenient to normalize separations with $\bar{\eta}$ and we will consider (10) at fixed $\ell^* = \ell / \bar{\eta}$.

In the limit of very small separations (i.e., $\ell^* \simeq 1$) (10) recovers the finite difference representation of the 1D-energy dissipation (7) and thus, from (8)

$$\frac{\langle E(\ell)^p \rangle}{\langle E(\ell) \rangle^p} = \frac{\langle \epsilon^p \rangle}{\langle \epsilon \rangle^p} \simeq \text{Re}_\lambda^{-2\theta_p}, \tag{12}$$

with the exponents θ_p given by (9).

In the case of separations in the inertial range, from (1) we have $\langle E(\ell)^p \rangle / \langle E(\ell) \rangle^p \simeq \ell^{\zeta_{2p} - p\zeta_2}$. Because $\ell = \ell^* \bar{\eta}$, using (11) we end with the prediction that the exponents of (12) are given by

$$\theta_p = \frac{\zeta_{2p} - p}{2 - \zeta_2}. \tag{13}$$

Let us remark that the set of exponents (9) and (13) are expected to be not very different (they are both zero for nonintermittent turbulence) and thus a discrimination between the two predictions require good accuracy. Scaling exponents (13) have been previously proposed to hold even in the dissipative scales.²¹ We will see that on these scales experimental data are more in favor of the set of exponents (9).

III. EXPERIMENTAL SET-UP AND DATA ANALYSIS

The experimental setup consists of a water jet in a closed circuit facility as shown in Fig. 1. A centrifugal pump moves water from a primary tank into a settling chamber which is




FIG. 1. Sketch of the experimental setup.

equipped with a valve to damp oscillations due to the pump. A series of contractions leads to a 1.5 m long pipe, with an inner diameter of 14 cm which is followed by the final contraction (1:50 in area) to the jet. The jet (diameter $d = 20$ mm) exits into a large water-filled tank (height $30d$, width $30d$, length $60d$) from which the water returns to the primary tank. The pipe, the contraction and large tank are made of perspex to allow optical access to the flow. At the jet exit, the flow is axisymmetric and has no swirl; preliminary measurements also confirmed that it is unaffected by any external forcing due to the pump. The jet has a top-hat velocity profile at the nozzle exit with a boundary layer shape parameter (defined as the ratio between the displacement and momentum thickness at the outlet) equal to 3.29 and a turbulent intensity equal to about 0.021 on the jet axis.

Velocity measurements are performed by means of a forward-scatter laser Doppler anemometer (LDA) equipped with two Bragg cells. The fringe spacing is $3.416 \mu\text{m}$ and the

measurement volume size is about 0.1, 0.1, and 0.8 mm along the x , y , and z axes, respectively. The LDA data are randomly distributed in time; therefore, they are resampled by using a linear interpolation to obtain evenly spaced samples and also to provide unbiased statistics.²² The value of Re_λ is sufficiently large to expect the LDA noise not to affect the behavior of structure functions in the IR.²³ The LDA data on longitudinal and transverse structure functions and on scaling exponents were compared to data obtained by hot wire anemometry in a similar jet; the agreement was good up to 8th order.²²

Measurements were made at $x/d \approx 40$ (x is measured from the nozzle exit plane), where the flow field may be considered to be approximately self-preserving and isotropic.²³ The jet exit velocity U_0 was selected so that the exit Reynolds number $Re_d \equiv U_0 d / \nu$ changes from about 2×10^4 to 2×10^5 . As a consequence, Re_λ changes from about 200 to almost 2000 at the measurement location. The number

TABLE I. Parameters of the experiments on the jet at $x/d = 40$ (except for the outlet mean velocity and the exit Reynolds number computed at $x/d = 0$).

Outlet mean velocity					
U_0 (cm/s)	102	196	304	487	1060
Re_d	20 400	39 200	60 800	97 400	21 2000
Local mean velocity	15.3	30.4	41.4	72.1	167
U (cm/s)					
Local rms velocity	4.4	9.2	12.3	23.8	50.1
u' (cm/s)					
Re_λ	230	308	435	926	1840
Local integral scale	5.5	6.3	9.4	22.0	41.1
L (cm)					
Local Kolmogorov scale	0.0181	0.0097	0.0086	0.0066	0.0045
η (cm)					




FIG. 2. Velocity power spectra at $Re_\lambda=230$ (thin line) and $Re_\lambda=1840$ (tick line) normalized with Kolmogorov scale and velocity. Dashed line represent the Kolmogorov spectrum, $E(k)=Ck^{-5/3}$ with $C=2$. In the inset we show in lin-log plot the compensated spectra $E(k)k^{5/3}$ together with the extrapolated spectrum at $Re_\lambda=230$ (dashed line) in order to shown the dissipative range.

of collected samples is about 10^6 in each run. As usually done in practice, we resort to Taylor hypothesis to transform time differences into space differences by means of the local mean velocity U . The number of independent samples, given by $UT_S/2L$ (where T_S is the total record duration), is about 10^4 . Probability density functions of the longitudinal velocity increments have been calculated at different values of ℓ^* . The distributions indicate that the number of samples is adequate for achieving a closure of the integrand associated with the structure functions $S_q(\ell)$ at least up to $q=6$. In Table I we summarize some parameters of the experiments.

Because the acquisition rate is constant for the different runs ($\delta t=5 \times 10^{-4}$ s) the smallest resolved scale $\delta x=U \delta t$ varies with Re_λ . By increasing Re_λ we have both an increasing of δx (as Re_λ^2) and a decreasing of the dissipative scale $\bar{\eta}$ according to (11). Together, they limit the possibility to completely resolve dissipative scales. In Fig. 2 we plot the velocity power spectra for the two extreme runs $Re_\lambda=230$ and $Re_\lambda=1840$ normalized with Kolmogorov scale and velocity.




FIG. 3. Structure function of order 2 and 4 for different values of Re_λ compensated with the scaling behavior (1). The x axis is adimensionalized with the Kolmogorov scale.

In the latter, more severe, case the compensated spectrum $E(k)k^{5/3}$ displays an exponential decay at large wave number up to about $k=0.15\eta^{-1}$. We can thus safely compute velocity differences down to $\ell^*\approx 10$ which is just at the border of the dissipative range.²⁴ Despite these limitations, we are confident that are able to capture, at least, the scaling behavior of small scale statistics, as shown below by the constancy of the mean energy dissipation.

In Fig. 3, the second and fourth-order structure functions obtained for $Re_\lambda=230$ and for $Re_\lambda=1840$ compensated with the scaling behavior (1) are plotted (Kolmogorov length and velocity scales are used to adimensionalize the horizontal and vertical axes). Scaling exponents can be derived from structure functions of Fig. 3 within the inertial range (approximately from $\ell\approx 80$ to $\ell\approx 500$, using a criterion based on 3% difference from the maximum in the third-order structure function divided by ℓ). The result ($\zeta_2=0.708\pm 0.020$ and $\zeta_4=1.265\pm 0.035$ for $Re_\lambda=230$) agrees with those obtained numerically and experimentally by other authors.⁴

IV. RESULTS AND COMPARISON WITH THEORETICAL PREDICTIONS

In Fig. 4 we plot the average surrogate energy dissipation $\langle E(\ell) \rangle$, computed from (10) and adimensionalized with large scale quantity u'^3/L , as a function of Re_λ . A constant value of energy dissipation is expected from energy balance arguments.³ Thus, the observed independence of $\langle E(\ell) \rangle$, computed at fixed $\ell^*=10$, on Re_λ is a confirmation that at least the scaling behavior of the dissipative scales is captured. We can give a different estimation of the dissipation on the basis of the energy spectrum. The advantage in this case is that we can reduce the noise by interpolating the spectrum at small scales, as shown in the inset of Fig. 2.^{3,25} The result for our data, also plotted in Fig. 4, is consistent with the previous one.

In order to verify the prediction for θ_p of Sec. II, the scaling laws (12) of the moments of $E(\ell)$ with Re_λ must be computed. In Fig. 5, an example of this scaling, computed at $\ell^*=10$, is given for different values of p . We are confident that at least up to $p=2$ the scaling exponents can be evaluated with an error on the fit less than 5%.




FIG. 4. Average energy dissipation computed from definition (10) (+) at $\ell^*=10$ and from the energy spectrum (\times) as function of Re_λ .




FIG. 5. Re_λ dependence of $\langle E(\ell)^p \rangle$ normalized with $\langle E(\ell) \rangle^p$ for $p=2/3$ (+), $p=4/3$ (x), and $p=2$ (*) at $\ell=10\eta$. The lines represent the best fit with a power law with exponent θ_p .

The resulting exponents θ_p are given in Fig. 6. They have been computed at two different values of the separation $\ell^*=10$ and $\ell^*=400$. As expected, for separations moving from the dissipative range into the inertial range, the curve of the exponents $\theta(p)$ becomes flatter and approaches the prediction (13). In Fig. 6 we also plot the two predictions (9) and (13). These have been obtained by computing the structure function scaling exponents ζ_q (1) as discussed above, at the largest Re_λ available. Given the exponents ζ_q , the codimension $Z(h)$ has been obtained by numerical inversion of (4). Finally, (9) is used to predict the values of θ_p . The agreement of our data with the two predictions (9) and (13) is remarkable. Some deviations are observed, especially for higher moments. Indeed, according to (10), the computation of θ_p corresponds roughly to the computation of the structure function of order $3p$.

The theoretical picture described in Sec. II is that the two sets of exponents (9) and (13) are asymptotically recovered for separations within the dissipative scales and the inertial range scales, respectively. This is indeed observed, as




FIG. 6. Exponents θ_p obtained from the data of Fig. 5 for $\ell=10\eta$ (+) and $\ell=400\eta$ (x). The continuous line is the intermittent prediction (9), the dashed line is the prediction (13). In the inset the value of θ_2 obtained at different scales is shown as a function of the scale.

shown in the inset of Fig. 6 where the value of θ_2 is plotted as a function of the separation. Despite the wide range of crossover, two constant asymptotic values are clearly observable.

Let us remark again that the difference between predictions (9) and (13) is based on the assumption of a dissipative scale fluctuating with the local energy dissipation. Our experimental data demonstrate that this is indeed the case.

V. REMARKS AND CONCLUSIONS

In this paper, the scaling of velocity increments at very small scales in a turbulent flow is investigated with special focus on Reynolds number dependence. Within the multifractal framework it is possible to derive a relation between the scaling of the energy dissipation statistics with the Reynolds number and the scaling exponents of the velocity structure functions at fixed Reynolds number. This relation is found to fit well experimental data taken into a water jet over about a decade of Re_λ . The one-dimensional surrogate of energy dissipation is estimated from velocity differences at different scales, from the border of the dissipative range to the deep inertial range. It is shown that our statistics gives asymptotically two sets of exponents for very small or inertial range separations. The rather wide region of crossover indicates a penetration of dissipative contribution within the inertial range.

The agreement of the experimental results with the two sets of exponents is a demonstration of the connection between dissipative scales and inertial range scales as described by multiplicative models. Moreover, the difference between the two sets of exponents is the signature of the fluctuating dissipative scale. If these results are dependent on the still finite value of the Reynolds number or on the anisotropy of the flow field is left as a subject for future investigations.

¹F. Anselmet, Y. Gagne, E. J. Hopfinger, and R. A. Antonia, "High-order velocity structure functions in turbulent shear flow," *J. Fluid Mech.* **140**, 63 (1984).
²A. Vincent and M. Meneguzzi, "The spatial structure and statistical properties of homogeneous turbulence," *J. Fluid Mech.* **225**, 1 (1991).
³A. S. Monin and A. M. Yaglom, *Statistical Fluid Mechanics* (MIT Press, Cambridge, MA, 1975).
⁴U. Frisch, *Turbulence* (Cambridge University Press, Cambridge, UK, 1995).
⁵R. A. Antonia, T. Zhou, and G. Xu, "Second-order temperature and velocity structure functions: Reynolds number dependence," *Phys. Fluids* **12**, 1509 (2000).
⁶R. A. Antonia, B. Pearson, and T. Zhou, "Reynolds number dependence of second-order velocity structure functions," *Phys. Fluids* **12**, 3000 (2000).
⁷A. Arneodo, C. Baudet, F. Belin, R. Benzi, B. Castaing, B. Chabaud, R. Chavarria, S. Ciliberto, R. Camussi, F. Chillà, B. Dubrulle, Y. Gagne, B. Hebral, J. Herweijer, M. Marchand, J. Maurer, J. F. Muzy, A. Naert, A. Noullez, J. Peinke, F. Roux, P. Tabeling, W. van de Water, and H. Willaime, "Structure functions in turbulence, in various flow configurations, at Reynolds number between 30 and 5000, using extended self-similarity," *Europhys. Lett.* **34**, 411 (1996).
⁸K. R. Sreenivasan, "On the scaling of the turbulence energy dissipation rate," *Phys. Fluids* **27**, 1048 (1984).
⁹K. R. Sreenivasan, "An update on the energy dissipation rate in isotropic turbulence," *Phys. Fluids* **10**, 528 (1998).
¹⁰C. Meneveau and K. R. Sreenivasan, "Simple multifractal cascade model for fully developed turbulence," *Phys. Rev. Lett.* **59**, 1424 (1987).
¹¹C. Meneveau and K. R. Sreenivasan, "The multifractal nature of turbulent energy dissipation," *J. Fluid Mech.* **224**, 429 (1991).

- ¹²R. Benzi, L. Biferale, G. Paladin, A. Vulpiani, and M. Vergassola, "Multifractality in the statistics of the velocity gradients in turbulence," *Phys. Rev. Lett.* **67**, 2299 (1991).
- ¹³K. Kailasnath, K. R. Sreenivasan, and G. Stolovitzky, "Probability density of velocity increments in turbulent flows," *Phys. Rev. Lett.* **68**, 2766 (1992).
- ¹⁴P. Tabeling, G. Zocchi, F. Belin, J. Maurer, and H. Willaime, "Probability density functions, skewness, and flatness in large Reynolds number turbulence," *Phys. Rev. E* **53**, 1613 (1996).
- ¹⁵G. Parisi and U. Frisch, "On the singularity structure of fully developed turbulence," in *Turbulence and Predictability in Geophysical Fluid Dynamics, Proceedings of the International School of Physics 'Environ. Fermi,' 1983, Varenna*, edited by M. Ghil, R. Benzi, and G. Parisi (North-Holland, Amsterdam, 1985).
- ¹⁶G. Paladin and A. Vulpiani, "Degrees of freedom of turbulence," *Phys. Rev. A* **35**, 1971 (1987).
- ¹⁷U. Frisch and M. Vergassola, "A prediction of the multifractal model: the intermediate dissipation range," *Europhys. Lett.* **14**, 439 (1991).
- ¹⁸C. Meneveau, "Transition between viscous and inertial-range scaling of turbulence structure functions," *Phys. Rev. E* **54**, 3657 (1996).
- ¹⁹M. Nelkin, "Multifractal scaling of velocity derivatives in turbulence," *Phys. Rev. A* **42**, 7226 (1990).
- ²⁰G. Boffetta, A. Celani, and D. Roagna, "Energy dissipation statistics in a shell model of turbulence," *Phys. Rev. E* **61**, 3234 (2000).
- ²¹J. Eggers and S. Grossmann, "Effect of dissipation fluctuations on anomalous velocity scaling in turbulence," *Phys. Rev. A* **45**, 2360 (1992).
- ²²G. P. Romano and R. A. Antonia, "Longitudinal and transverse structure functions in a turbulent round jet: effect of initial conditions and Reynolds number," *J. Fluid Mech.* **436**, 231 (2001).
- ²³R. A. Antonia, T. Zhou, and G. P. Romano, "Second- and third-order longitudinal velocity structure functions in a fully developed turbulent channel flow," *Phys. Fluids* **9**, 3465 (1997).
- ²⁴R. A. Antonia and B. Pearson, "Effect of initial conditions on the mean energy dissipation rate and the scaling exponent," *Phys. Rev. E* **62**, 8086 (2000).
- ²⁵R. A. Antonia and B. Pearson, "Reynolds number dependence of velocity structure functions in a turbulent pipe flow," *Flow, Turbul. Combust.* **64**, 95 (2000).