

Split energy cascade in turbulent thin fluid layers

Stefano Musacchio and , and Guido Boffetta

Citation: *Physics of Fluids* **29**, 111106 (2017); doi: 10.1063/1.4986001

View online: <http://dx.doi.org/10.1063/1.4986001>

View Table of Contents: <http://aip.scitation.org/toc/phf/29/11>

Published by the *American Institute of Physics*

**COMPLETELY
REDESIGNED!**



**PHYSICS
TODAY**

Physics Today Buyer's Guide
Search with a purpose.

Split energy cascade in turbulent thin fluid layers

Stefano Musacchio¹ and Guido Boffetta²

¹Université Côte d'Azur, CNRS, LJAD, Nice, France

²Dipartimento di Fisica and INFN, Università di Torino, Via P. Giuria 1, 10125 Torino, Italy

(Received 31 May 2017; accepted 24 July 2017; published online 9 August 2017)

We discuss the phenomenology of the split energy cascade in a three-dimensional thin fluid layer by means of high resolution numerical simulations of the Navier-Stokes equations. We observe the presence of both an inverse energy cascade at large scales, as predicted for two-dimensional turbulence, and a direct energy cascade at small scales, as in three-dimensional turbulence. The inverse energy cascade is associated with a direct cascade of enstrophy in the intermediate range of scales. Notably, we find that the inverse cascade of energy in this system is not a purely 2D phenomenon, as the coupling with the 3D velocity field is necessary to guarantee the constancy of fluxes. *Published by AIP Publishing.* [<http://dx.doi.org/10.1063/1.4986001>]

I. INTRODUCTION

Fifty years ago, Kraichnan showed in his seminal paper¹ that the dynamics of an incompressible flow in two-dimensions (2D) are dramatically different from the classical phenomenology of three-dimensional (3D) turbulence. The presence of two inviscid quadratic invariants, energy and enstrophy, gives rise to a double-cascade scenario.² At variance with the 3D case, in which the kinetic energy cascades toward small viscous scales, in 2D it is transferred toward large scales. Such an “inverse energy cascade” is accompanied by a “direct enstrophy cascade,” which proceeds towards small scales.³ In the inverse and direct ranges of scales, the theory predicts a kinetic energy spectrum $E(k) \simeq \varepsilon_f^{2/3} k^{-5/3}$ and $E(k) \simeq \eta_f^{2/3} k^{-3}$ with possible logarithmic corrections.⁴ Here and in the following, ε_f and η_f denote the energy and the enstrophy injection rates, respectively. Kraichnan's seminal concept of inverse cascade has since become a prototypical model for several turbulent systems, from the inverse cascade in strongly rotating 3D flows⁵ to the inverse cascade of magnetic helicity in three-dimensional magneto-hydrodynamic turbulence,⁶ of passive scalar in compressible turbulence,⁷ and of wave action in weak turbulence.⁸

The presence of the two cascades in two-dimensional turbulence has been observed in a number of numerical simulations^{9–18} and in experiments on soap films^{19–25} and in thin fluid layers.^{26–28}

At variance with the numerical investigations, which allow the study of the ideal 2D Navier-Stokes equations, the experiments have to deal with the effects of the finite thickness of the fluid layer. This issue, which has been often considered a limitation for two-dimensional experiments, opens a series of interesting questions. How is the ideal 2D phenomenology modified in the case of a thin (3D) fluid layer? How thin should the layer be, in order to display the 2D-like double cascade? In which way does the transition from the 2D to the 3D regime occur at increasing the thickness of the layer?

These questions have been addressed both in numerical simulations^{29,30} and in experiments,^{31,32} which have shown

that the critical parameter that controls the cascade direction is the ratio $S = L_z/L_f$ of the thickness to the forcing scale. When $S \geq 1$, the flow is 3D at the forcing scale and the injected energy produces a direct cascade as in usual 3D turbulence. By decreasing S , one observes the phenomenon of *cascade splitting*, with a fraction of the injected energy going to large scales and the remaining energy flowing to small scales.^{29,30} For $S \ll 1$, the flux of the direct energy cascade is expected to scale as S^2 . This prediction has been verified in shell models for quasi-two-dimensional turbulence.³³ Further reducing S , the cascade of energy towards small scales vanishes when the thickness L_z reaches the Kolmogorov scale L_ν and the flow recovers the standard 2D phenomenology.

In the present paper, we investigate the entanglement of 2D and 3D dynamics that occurs in a turbulent fluid layer by means of numerical simulations of the 3D Navier-Stokes equations in a confined domain with $L_z < L_f$ (and $L_z > L_\nu$). In agreement with previous findings, we observe the phenomenon of splitting of the energy cascade. By introducing a suitable decomposition of the velocity field, we show that the inverse cascade mainly involves the kinetic energy of the 2D modes, while the energy of the remnant 3D velocity is transferred toward the viscous scales. We also show that the development of the inverse energy cascade is associated with a partial conservation of the enstrophy in the intermediate range of scales $L_z < \ell < L_f$. Interestingly, we find that 3D modes play a relevant role in the 2D phenomenology which is observed at large scales. In particular, the transport of the 2D modes by the 3D velocity is necessary to ensure a constant flux of energy in the inverse cascade as well as a constant flux of enstrophy in the range $L_z < \ell < L_f$.

The remaining part of the paper is organized as follows. Section II introduces the Navier-Stokes equations and the decomposition of the velocity field in the 2D and 3D modes. In Sec. III, we report the results of the numerical simulations. Section IV is devoted to the conclusions. In the Appendix, we derive a 2D model for the dynamics of a thin layer in the limit $L_z \rightarrow 0$.

II. NAVIER-STOKES EQUATIONS FOR A THIN LAYER

We consider the dynamics of a three-dimensional thin layer of fluid, ruled by the Navier-Stokes equations for the velocity field $\mathbf{u}(\mathbf{x}, t)$,

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u} + \mathbf{f}, \quad (1)$$

where ν is the kinematic viscosity, \mathbf{f} is the external forcing, and the pressure p is determined by the incompressibility constraint $\nabla \cdot \mathbf{u} = 0$. The flow is confined in a thin domain of size $L_x = L_y = rL_z$ with $r \gg 1$ and periodic boundary conditions in all the directions.

In the absence of forcing and dissipation, (1) preserves the kinetic energy $E = (1/2)\langle |\mathbf{u}|^2 \rangle$ (where $\langle \dots \rangle$ denotes average over the space). The energy balance in the forced-dissipated case reads

$$\frac{dE}{dt} = \varepsilon_f - \varepsilon_\nu, \quad (2)$$

where $\varepsilon_\nu = \nu \langle (\nabla \mathbf{u})^2 \rangle$ is the energy dissipation rate due to the viscosity and $\varepsilon_f = \langle \mathbf{f} \cdot \mathbf{u} \rangle$ is the energy input provided by the external forcing.

The forcing also provides an input of enstrophy $Z = (1/2)\langle |\boldsymbol{\omega}|^2 \rangle$, ($\boldsymbol{\omega} = \nabla \times \mathbf{u}$ denotes the vorticity field) at the rate $\eta_f = \langle (\nabla \times \mathbf{f}) \cdot \boldsymbol{\omega} \rangle$. At variance with the ideal 2D case, the enstrophy is not preserved by the fully 3D inviscid dynamics, but it is produced by the vortex stretching mechanism.³⁴ The equation for the vorticity field is obtained by taking the curl of (1),

$$\partial_t \boldsymbol{\omega} + \mathbf{u} \cdot \nabla \boldsymbol{\omega} = \boldsymbol{\omega} \cdot \nabla \mathbf{u} + \nu \Delta \boldsymbol{\omega} + \mathbf{f}_\omega, \quad (3)$$

where $\mathbf{f}_\omega = \nabla \times \mathbf{f}$ and $\boldsymbol{\omega} \cdot \nabla \mathbf{u}$ represents the vortex stretching term.

In order to highlight the presence of a 2D phenomenology in the 3D flow, it is useful to decompose the velocity field as $\mathbf{u} = \mathbf{u}^{2D} + \mathbf{u}^{3D}$. The 2D mode $\mathbf{u}^{2D} = (u_x^{2D}(x, y), u_y^{2D}(x, y), 0)$ is defined as the average along the z direction of the x and y components of the velocity field \mathbf{u} , and it satisfies the 2D incompressibility condition $\partial_x u_x^{2D} + \partial_y u_y^{2D} = 0$. In the Fourier space, it corresponds to the mode $k_3 = 0$ of the horizontal velocity $\mathbf{u}^{2D}(k_1, k_2) = (u_x(k_1, k_2), u_y(k_1, k_2), 0)$. The field \mathbf{u}^{3D} is defined as the difference $\mathbf{u}^{3D} = \mathbf{u} - \mathbf{u}^{2D}$. By the above definitions, it is easy to show that the total energy decomposes into a 2D contribution $E^{2D} = (1/2)\langle |\mathbf{u}^{2D}|^2 \rangle$ and a 3D contribution $E^{3D} = (1/2)\langle |\mathbf{u}^{3D}|^2 \rangle$ as $E = E^{2D} + E^{3D}$. We notice that, beside the kinetic energy of the vertical velocity, E^{3D} also contains the contributions of the modes $k_3 \neq 0$ of the horizontal components of the velocity.

Similarly, the vorticity field can be decomposed as $\boldsymbol{\omega} = \omega^{2D} \hat{\mathbf{z}} + \boldsymbol{\omega}^{3D}$, where $\omega^{2D} = \partial_x u_y^{2D} - \partial_y u_x^{2D}$ is the scalar vorticity of the two-dimensional flow. In the limit of vanishing thickness $L_z \rightarrow 0$ (at finite viscosity ν), the vertical dependence disappears and \mathbf{u}^{2D} becomes solution of the 2D Navier-Stokes equation (see the Appendix). Therefore, it is reasonable to assume that the occurrence of an inverse energy cascade at a finite thickness L_z should be associated with the dynamics of the 2D mode \mathbf{u}^{2D} .

III. DIRECT NUMERICAL SIMULATIONS

We performed a direct numerical simulation of the Navier-Stokes equations (1) in a confined geometry with periodic

boundary conditions. The computational domain has dimensions $L_x = L_y = 2\pi$, $L_z = L_x/64$ ($r = 64$), and it is discretized on a uniform grid at the resolution $N_x \times N_y \times N_z = 4096 \times 4096 \times 64$. The numerical simulations are performed by means of a fully parallel, pseudospectral code, with a 2/3 dealiasing scheme. We adopt a hyperviscous damping scheme $(-1)^{p-1} \nu_p \Delta^p$ with $p = 4$ and $\nu_p = 10^{-21}$. We do not use any large-scale dissipation (such as linear friction).

The flow is sustained by a ‘‘two-components, two-dimensional’’ forcing; that is, the forcing is active on the horizontal components of the velocity, and it is dependent on the horizontal coordinates only, $\mathbf{f} = (f_x(x, y), f_y(x, y), 0)$. Therefore, in the vorticity equation (3), the forcing \mathbf{f}_ω is active only on the 2D field ω^{2D} . The forcing is restricted to a narrow wavenumber shell in the Fourier space with $k_h = (k_1^2 + k_2^2)^{1/2} \simeq k_f$, $k_3 = 0$. Here $k_f = 16$. The forcing is Gaussian and δ -correlated in time to control the injection rates ε_f and η_f . The characteristic time at the forcing scale is defined as $\tau_f = \eta_f^{-1/3}$. The ratio between the thickness $L_z = 2\pi/k_z$ and the forcing scale $L_f = 2\pi/k_f$ is $S = L_z/L_f = 4$. This ensures the regime of split cascade with the coexistence of the 3D and 2D phenomenologies (see Fig. 1).³⁰

The velocity field at time $t = 0$ is initialized to zero plus a small random perturbation that is required to trigger the 3D instability. The results shown in this section have been obtained with an energy of the initial perturbation $E_{pert} \simeq 1.6 \times 10^{-7} \varepsilon_f \tau_f$.

Because of the purely 2D forcing, in the early stage of the simulation ($t < 10\tau_f$), the energy accumulates in the 2D mode \mathbf{u}^{2D} at a rate equal to the forcing input, while the energy E^{3D} of the 3D component is negligible (see Fig. 2). The activation of the 3D modes \mathbf{u}^{3D} (at $t \simeq 10\tau_f$) is accompanied by a significant reduction of the growth rate of the 2D energy as some of the injected energy is now transferred to small scales. At later times, E^{3D} saturates to a statistically steady value (as in standard 3D turbulence), while the two-dimensional component E^{2D} energy keeps increasing with a constant reduced rate $\varepsilon_\alpha < \varepsilon_f$. We stop the simulation at time $t = 100\tau_f$, when the inverse energy cascade has reached the lowest wavenumber (see Fig. 4). Continuing the simulation further, in the absence




FIG. 1. Snapshot of the scalar vorticity field ω^{2D} in the late stage of the simulation. Typical two-dimensional objects, such as strong vortices at the forcing scale L_f , coexist with small scale three-dimensional features.




FIG. 2. Temporal evolution of the kinetic energy E^{2D} (red, dashed line, left y-axis) and E^{3D} (blue, solid line, right y-axis). The dotted line represents the linear growth with the input rate ϵ_f (left y-axis).

of large-scale dissipation, we expect that the kinetic energy will accumulate in the lowest mode, giving rise to the so called condensate.^{35–37} The 2D mode contains almost all the kinetic energy of the horizontal velocities, and the kinetic energy of the vertical component u_z contained in the mode $k_3 = 0$ represents only 24% of the total.

The enstrophy of the 2D mode $Z^{2D} = 1/2\langle(\omega^{2D})^2\rangle$ grows initially with the input rate η_f , reflecting the purely 2D nature of the initial flow (see Fig. 3). As 3D motions develop (at $t \simeq 10\tau_f$), the enstrophy associated with the 3D modes $Z^{3D} = 1/2\langle|\omega^{3D}|^2\rangle$ increases very rapidly and reaches a stationary value which is much larger than the saturation level of Z^{2D} .

A. Energy spectra

In Fig. 4, we show the instantaneous spectra $E(k, t)$ of the total energy at different times t . The initial spectra are almost completely 2D because the forcing is active only on 2D modes. We observe that the 3D instability begins at high harmonics of the thickness wavenumber k_z , then propagating to all the modes $k > k_z$. The energy spectrum at $k > k_f$ saturates at time $t \simeq 16\tau_f$. At scales larger than the forcing scale, $k < k_f$, we




FIG. 3. Temporal evolution of the enstrophy of ω^{2D} (red, dashed line, left y-axis) and ω^{3D} (blue, solid line, right y-axis). The dotted line represents the linear growth with the input rate η_f (left y-axis).




FIG. 4. Energy spectra $E(k, t)$ at times $t/\tau_f = 4.6$ (a, violet), 6.9 (b, green), 9.2 (c, cyan), 13.7 (d, orange), 18.3 (e, yellow), 45.8 (f, blue), and 91.6 (g, red). The last three spectra are almost superposed for $k > k_z$.

observe the development of an inverse energy cascade with a power-law spectrum $E(k) \simeq \epsilon_\alpha^{2/3} k^{-5/3}$.

The “spiky” aspect of the energy spectrum at high wavenumbers $k > k_z$ is due to the anisotropic spacing of the wavenumbers in the Fourier space. The separation between the discrete wavenumbers in the horizontal direction is $\Delta k_h = 2\pi/L_x$, while in the vertical direction it is $\Delta k_3 = 2\pi/L_z = k_z$. Given that $L_z \ll L_x$, the wavenumber space is structured as horizontal dense layers, separated by large gaps in the vertical direction. Because the complete energy spectrum is defined as the integral of the square amplitude of the modes over a spherical wavenumber shell of radius k , one gets a sudden increase of the spectrum each time the spherical shell is a multiple of k_z .

In order to analyze the contribution of the 2D mode to the total energy spectrum, it is also interesting to consider the 2D spectra $E^{2D}(k)$, in which the integral is restricted to the horizontal wavenumbers $k_h = \sqrt{k_1^2 + k_2^2}$ on the plane $k_3 = 0$. In physical space, this is equivalent to averaging the velocity fields in the vertical direction z first and then computing the spectrum of the averaged 2D fields. It is worth to notice that for $k < k_z$, the 2D spectra and 3D spectra coincide because the spherical shell of radius $k < k_z$ intersects the planes $k_3 = \pm mk_z$ only for $m = 0$.

In Fig. 5, we compare the 2D energy spectrum $E^{2D}(k)$ of the 2D mode \mathbf{u}^{2D} with the 3D energy spectrum of the 3D mode \mathbf{u}^{3D} . At low wavenumbers $k < k_z$, almost all the kinetic energy is contained in the 2D mode. This confirms that the inverse cascade that develops in the range $k < k_f$ concerns only the 2D energy. Conversely, the 3D mode contains the largest fraction of the kinetic energy at high wavenumbers $k > k_z$. Interestingly, in the same range of wavenumbers, the 2D spectrum of the 2D mode displays a $-5/3$ slope, and it is very close to the 2D spectrum of the vertical component u_z .

The spectrum $E^{2D}(k)$ shown in Fig. 5 is reminiscent of the horizontal spectrum (of meridional and zonal winds) observed in the upper troposphere by the Global Atmospheric Sampling Program³⁸ where a transition from a k^{-3} to a $k^{-5/3}$ spectrum at small scales is observed. We remark that despite the similarities between the two spectra, the physical mechanisms are




FIG. 5. Energy spectrum $E^{2D}(k)$ of the 2D mode u^{2D} (red, dashed line) and 3D mode u^{3D} (blue, solid line). We also show the 2D spectrum of the vertically averaged vertical velocity u_z (black, dotted line). The spectra are computed at $t = 100\tau_f$.

probably different (see, for example, Refs. 39 and 40) as the transition in the atmospheric spectrum is observed at a scale around 500 km, while in our simulations it occurs at a scale comparable with the thickness of the layer.

B. Spectral fluxes

The analysis of the spectral flux of the total kinetic energy $\Pi_E(k)$ shows that in a thin fluid layer, the energy injected by the forcing at the wavenumber k_f indeed splits into two parts (see Fig. 6). At low wavenumbers $k < k_f$, we observe a constant, negative flux of energy, which indicates the presence of an inverse energy transfer toward large scales. At high wavenumbers $k > k_f$, we also observe a constant energy flux, now positive, indicating a direct cascade of energy toward small scales. It is worth reminding that the kinetic energy transported in the direct cascade is not only that of the vertical component of the velocity but contains also the contributions of the modes $k_3 \neq 0$ of the horizontal velocities. In this range of scales, the dynamics in the vertical and horizontal directions are strongly coupled, and the positive energy flux cannot be explained in terms of a direct cascade of the vertical velocity passively transported by a two-dimensional, three components (2D3C) flow.^{41–43}




FIG. 6. Spectral energy flux $\Pi_E(k)$ (blue solid line) and 2D energy flux (red, dashed line), at $t = 100\tau_f$.

As we have shown in Fig. 4, the inverse cascade mainly involves the energy of the 2D mode. This observation suggests one to check whether or not this inverse cascade coincides with a purely 2D dynamics of the 2D mode u^{2D} . To this purpose, we have taken the fields u^{2D} and truncated them at $k_h = k_z$ by setting to zero all the modes with $k_h > k_z$. Then, we have computed the 2D spectral fluxes of the truncated 2D fields, assuming that they were solutions of the two-dimensional Navier-Stokes equations. Surprisingly, we find that the 2D energy flux in the range of scales of the inverse cascade does not coincide with the 3D flux shown in Fig. 5. The physical interpretation of this result is that the energy of the 2D mode is not simply transported toward large scales by the 2D flow itself, but the 3D modes contribute to the transport process. This contrasts with a 2D3C scenario at large scales, with the vertical velocity being passively transported by the 2D flow.^{41–43}

As discussed in Sec. II, the main difference between the 3D and 2D Navier-Stokes equations is the absence of the vortex stretching term $\omega \cdot \nabla u$ in the latter. The presence of two positive-defined inviscid invariants (energy and enstrophy) causes the reversal of the direction of the energy cascade. Even if the enstrophy is not conserved by the 3D dynamics, it is tempting to conjecture that the development of the inverse cascade in thin fluid layers is due to a dynamical suppression of the enstrophy production. To investigate this issue, we computed the total spectral enstrophy flux $\Pi_Z(k)$ and the total spectral enstrophy production $\Sigma_Z(k)$ defined as

$$\Pi_Z = \int_{|q| \leq k} d\mathbf{q} (\mathbf{v} \cdot \nabla \omega)(\mathbf{q}) \omega^*(\mathbf{q}), \quad (4)$$

$$\Sigma_Z = \int_{|q| \leq k} d\mathbf{q} (\omega \cdot \nabla \mathbf{v})(\mathbf{q}) \omega^*(\mathbf{q}). \quad (5)$$

In the range of wavenumbers $k_f < k < k_z$, the production of enstrophy is negligible and the enstrophy flux is constant, as shown in Fig. 7. This constant flux corresponds to the presence of a direct enstrophy cascade. At high wavenumbers, $k > k_z$, the enstrophy production becomes significant and, therefore, the enstrophy flux is not constant anymore but grows following the production term. Our results show that in analogy with the




FIG. 7. Spectral enstrophy flux $\Pi_Z(k)$ (red solid line), spectral enstrophy production $\Sigma_Z(k)$ (blue, dashed line), and 2D enstrophy flux (black, dotted line), at $t = 100\tau_f$.

case of ideal 2D Navier-Stokes equations, also in a thin fluid layer, the emergence of the inverse energy cascade is due to the presence of a “quasi invariant,” the enstrophy, which is conserved by the large-scale dynamics. Nonetheless, it is worth to notice that the conservation of the enstrophy is not due to the transport by the 2D mode itself. Following the same procedure described in the case of the energy flux, we have computed the 2D enstrophy flux from the truncated 2D velocity fields. In the range of scale $k_f < k < k_z$, the 2D flux is positive, but not constant, thus indicating that the fully 3D dynamics are required for the conservation of enstrophy (see Fig. 7).

The presence of an intermediate direct enstrophy cascade in the range of scales $L_z < \ell < L_f$ allows deriving a simple dimensional argument for the scaling of the energy flux toward small scales. A direct cascade with a constant enstrophy flux $\eta_f = \varepsilon_f k_f^2$ also carries a residual energy flux, which decreases as $\Pi(k) \sim \varepsilon_f (k_f/k)^2$. By assuming that the flux of the energy of the direct cascade which starts from k_z is equal to the residual flux $\Pi(k_z)$ carried by the enstrophy cascade at the scale k_z , one gets the prediction $\varepsilon_v = \Pi(k_z) \sim \varepsilon_f S^2$. This prediction has been verified in shell models for quasi-two-dimensional turbulence.³³

IV. CONCLUSIONS

In this paper we present a numerical study of the phenomenology of a turbulent flow confined in a thin fluid layer. We discuss the possibility to disentangle the complex mixture of 2D and 3D dynamics by a suitable decomposition of the velocity field in 2D and 3D modes.

By analogy with previous studies,^{29,30} when the flow is forced at scales L_f larger than the thickness L_z of the layer, we observe a splitting of the energy cascade in two directions. A fraction of the energy is transported toward large scales, giving rise to an inverse energy cascade, while the remnant energy is transported toward the small viscous scales, as in 3D turbulence. We show that the inverse energy cascade is accompanied by the development of a direct cascade of enstrophy in the intermediate range of scales $L_z < \ell < L_f$. The enstrophy production becomes relevant only at small scales $\ell < L_z$, allowing for a partial conservation of the enstrophy by the large-scale dynamics.

The scenario that emerges from our findings is a coexistence of 2D phenomenology, with a double cascade of energy and enstrophy *à la Kraichnan* at large scales $\ell > L_z$ and a 3D direct energy cascade *à la Kolmogorov* at small scales $\ell < L_z$. Interestingly, the decomposition of the velocity field in the 2D modes and the remaining 3D part reveals that the 2D and 3D dynamics are deeply entangled. On the one hand, we find that the energy and enstrophy that are involved in the double cascade at large scales are those of the 2D modes. On the other hand, the 3D velocity is necessary to guarantee a constant flux of 2D energy and enstrophy in the large-scale transport.

We plan to extend the analysis of the interactions between 2D and 3D modes to the case of rotating and stably stratified thin fluid layers. Previous results⁴⁴ show that rotation causes a suppression of the enstrophy production similar to the effects of confinement, favoring the two-dimensionalization of

the flow and the development of the inverse energy cascade. Nonetheless, this effect is not accompanied by the presence of a range of scales in which the enstrophy is conserved by the large-scale dynamics. This is likely to affect the interactions between the 2D and 3D modes. In the case of stably stratified fluid layers, it has been shown that the conversion of kinetic energy into potential energy, which is promptly transferred toward the small diffusive scales, provides a fast dissipative mechanism that suppresses the large scale energy transfer.⁴⁵ Investigating the interactions between 2D vortical modes and 3D potential modes will improve the understanding of this process.

ACKNOWLEDGMENTS

Simulations have been performed at the Juelich Forschungszentrum (Germany) within the PRACE Preparatory Project No. PRPA22 and at CSC (Finland) within the European project HPC-Europa2 “Energy transfer in turbulent fluid layers.”

We acknowledge the useful discussions with A. Celani and P. Muratore Ginanneschi and the support by the European COST Action MP1305 “Flowing Matter.”

APPENDIX: 2D MODEL FOR THIN LAYERS

In this Appendix we discuss a two-dimensional model to describe the dynamics of a thin layer in which both the thickness L_z and the Kolmogorov scale L_v tend to zero, but their ratio remains of order unity.

We consider the Navier-Stokes equations (1) in a box in which the horizontal dimensions $L_x = L_y = L$ are much larger than the vertical thickness $L_z = \epsilon L$. The aspect ratio of the box is determined by the ratio $\epsilon = L_x/L_z$. We assume periodic boundary conditions in all the directions. We assume also that the external force \mathbf{f} acts only on the horizontal components and depends only on the horizontal coordinates: $\mathbf{f}(\mathbf{x}) = (f_x(x, y), f_y(x, y), 0)$.

The assumptions that the thickness L_z of the layer is very small and that it is of the order of the viscous scale allows one to suppose that the modes $k_3 > k_z$ are suppressed by the viscosity and can be neglected. Therefore, we make a Fourier truncation in the vertical direction by retaining only the first modes in the vertical direction $k_3 = 0, \pm k_z$, where $k_z = 2\pi/L_z = O(\epsilon^{-1})$. The velocity fields can be expanded as

$$u_l = u_l^0 + \sqrt{2} \left[u_l^c \cos(k_z z) + u_l^s \sin(k_z z) \right] \quad (A1)$$

$$u_3 = u_3^0 + \sqrt{2} \left[u_3^c \cos(k_z z) + u_3^s \sin(k_z z) \right], \quad (A2)$$

where $l \in [1, 2]$. Within this notation, the 2D mode \mathbf{u}^{2D} introduced in Sec. II coincides with the field $\mathbf{u}^0 = (u_1^0, u_2^0)$.

The incompressibility condition $\nabla \cdot \mathbf{u} = 0$ gives

$$\partial_l u_l^0 = 0; \quad \partial_l u_l^c = -k_z u_3^s; \quad \partial_l u_l^s = k_z u_3^c. \quad (A3)$$

This shows that the 2D mode \mathbf{u}^0 satisfies the 2D incompressibility. The fields $u_3^{s,c}$ are determined by the compressibility of the fields $u_l^{s,c}$. Expanding at leading order in ϵ the Navier-Stokes equations (1), one obtains the equations for the fields

u_l^0 , $u_l^{s,c}$, and u_3^0 ,

$$\partial_t u_l^0 + u_n^0 \partial_n u_l^0 = -\partial_l p + \nu \partial_n \partial_n u_l^0 + f_l \quad (\text{A4})$$

$$- \partial_n (u_n^c u_l^c + u_n^s u_l^s),$$

$$\partial_t u_l^c + u_n^0 \partial_n u_l^c = \nu \partial_n \partial_n u_l^c - \alpha u_l^c \quad (\text{A5})$$

$$- u_n^c \partial_n u_l^0 + u_3^0 k_z u_l^s,$$

$$\partial_t u_l^s + u_n^0 \partial_n u_l^s = \nu \partial_n \partial_n u_l^s - \alpha u_l^s \quad (\text{A6})$$

$$- u_n^s \partial_n u_l^0 - u_3^0 k_z u_l^c,$$

$$\partial_t u_3^0 + u_n^0 \partial_n u_3^0 = \nu \partial_n \partial_n u_3^0 \quad (\text{A7})$$

$$+ (u_n^c \partial_n \partial_l u_l^s - u_n^s \partial_n \partial_l u_l^c) / k_z,$$

where $l, n \in [1, 2]$ and summation over repeated indices is assumed.

The linear friction term $-\alpha \mathbf{u}^{s,c}$ in the 2D model comes from the derivatives in the vertical direction of the viscous term in Eq. (1). At finite viscosity ν , in the limit $k_z \rightarrow \infty$, the friction coefficient $\alpha = \nu k_z^2$ diverges. The modes $\mathbf{u}^{c,s}$ are, therefore, exponentially suppressed. In this limit, the equation for the 2D mode \mathbf{u}^0 reduces to the two-dimensional Navier-Stokes equations.

It is also interesting to consider the limit $k_z \rightarrow \infty$, in which the viscosity vanishes as $\nu \sim k_z^{-2} \rightarrow 0$ such that $\alpha = \nu k_z^2 \rightarrow \text{const.}$ In this case, the 2D mode \mathbf{u}^0 remains coupled with the 3D modes $\mathbf{u}^{s,c}$. The latter are transported and stretched by the gradients of the velocity field \mathbf{u}^0 and are coupled among themselves as a harmonic oscillator whose frequency is determined by the scalar field u_3^0 .

The stretching of the fields $\mathbf{u}^{s,c}$ is contrasted by the linear relaxation term $-\alpha \mathbf{u}^{s,c}$. If the dissipation prevails, the two fields are exponentially damped and their feedback on the 2D velocity field \mathbf{u}^0 can be neglected. Conversely, when stretching dominates, part of the kinetic energy is transferred to the fields $\mathbf{u}^{s,c}$ and is dissipated by the friction. The transition between the two regimes is expected to occur when $\lambda \sim \alpha$, where λ is a suitable measure of the intensity of the gradients of the 2D mode. A dimensional estimate based on the scaling laws of the direct enstrophy cascade gives $\lambda \sim \eta_f^{1/3}$, where η_f is the enstrophy flux. Recalling that the viscous scale L_ν (defined as the scale at which $Re = 1$) is $L_\nu = \nu^{1/2} \eta_f^{-1/6}$ and that $\alpha \sim \nu L_z^{-2}$, the condition $\lambda > \alpha$ for the transition is equivalent to $L_z > L_\nu$. This shows that 3D modes can be excited only when the thickness L_z is larger than the viscous scale L_ν .

The development of the 3D modes causes a reduction of the growth rate of the 2D mode. This can be seen from the energy balance of the 2D model

$$\frac{d}{dt} (E^{2D} + E^{3D}) = \varepsilon_f - 2\alpha E^{3D}, \quad (\text{A8})$$

where $E^{2D} = 1/2 \langle |\mathbf{u}^0|^2 \rangle$ is the energy of the 2D mode and $E^{3D} = 1/2 \langle |\mathbf{u}^c|^2 + |\mathbf{u}^s|^2 \rangle$ is the energy of the 3D modes. For $L_z > L_\nu$, the E^{3D} attains a positive, statistically constant value, and therefore the growth rate dE^{2D}/dt reduces to $\varepsilon_\alpha = \varepsilon_f - 2\alpha E^{3D} < \varepsilon_f$.

We notice that the model does not provide a quantitative estimate of the energy E^{3D} , which appears in Eq. (A8), and

therefore it is not possible to determine the scaling dependence of the growth rate of E^{2D} on the ratio L_z/L_ν .

- ¹R. H. Kraichnan, "Inertial ranges in two-dimensional turbulence," *Phys. Fluids* **10**, 1417 (1967).
- ²G. L. Eyink, "Exact results on stationary turbulence in 2D: Consequences of vorticity conservation," *Phys. D* **91**, 97 (1996).
- ³G. Boffetta and R. E. Ecke, "Two-dimensional turbulence," *Annu. Rev. Fluid Mech.* **44**, 427 (2012).
- ⁴R. H. Kraichnan, "Inertial-range transfer in two- and three-dimensional turbulence," *J. Fluid Mech.* **47**, 525 (1971).
- ⁵A. Pouquet, A. Sen, D. Rosenberg, P. D. Mininni, and J. Baerenzung, "Inverse cascades in turbulence and the case of rotating flows," *Phys. Scr.* **2013**, 014032.
- ⁶U. Frisch, A. Pouquet, J. Léorat, and A. Mazure, "Possibility of an inverse cascade of magnetic helicity in magnetohydrodynamic turbulence," *J. Fluid Mech.* **68**, 769 (1975).
- ⁷M. Chertkov, I. Kolokolov, and M. Vergassola, "Inverse versus direct cascades in turbulent advection," *Phys. Rev. Lett.* **80**, 512 (1998).
- ⁸V. E. Zakharov and M. M. Zaslavskii, "The kinetic equation and Kolmogorov spectra in the weak turbulence theory of wind waves," *Izv. Atmos. Oceanic Phys.* **18**, 747 (1982).
- ⁹U. Frisch and P.-L. Sulem, "Numerical simulation of the inverse cascade in two-dimensional turbulence," *Phys. Fluids* **27**, 1921 (1984).
- ¹⁰J. R. Herring and J. C. McWilliams, "Comparison of direct numerical simulation of two-dimensional turbulence with two-point closure: The effects of intermittency," *J. Fluid Mech.* **153**, 229 (1985).
- ¹¹M. E. Maltrud and G.K. Vallis, "Energy spectra and coherent structures in forced two-dimensional and beta-plane turbulence," *J. Fluid Mech.* **228**, 321 (1991).
- ¹²L. M. Smith and V. Yakhot, "Bose condensation and small-scale structure generation in a random force driven 2D turbulence," *Phys. Rev. Lett.* **71**, 352 (1993).
- ¹³G. Boffetta, A. Celani, and M. Vergassola, "Inverse energy cascade in two-dimensional turbulence: Deviations from Gaussian behavior," *Phys. Rev. E* **61**, R29 (2000).
- ¹⁴G. Boffetta, A. Celani, S. Musacchio, and M. Vergassola, "Intermittency in two-dimensional Ekman-Navier-Stokes turbulence," *Phys. Rev. E* **66**, 026304 (2002).
- ¹⁵S. Chen, R. E. Ecke, G. L. Eyink, X. Wang, and Z. Xiao, "Physical mechanism of the two-dimensional enstrophy cascade," *Phys. Rev. Lett.* **91**, 214501 (2003).
- ¹⁶Z. Xiao, M. Wan, S. Chen, and G. L. Eyink, "Physical mechanism of the inverse energy cascade of two-dimensional turbulence: A numerical investigation," *J. Fluid Mech.* **619**, 1 (2009).
- ¹⁷G. Boffetta and S. Musacchio, "Evidence for the double cascade scenario in two-dimensional turbulence," *Phys. Rev. E* **82**, 016307 (2010).
- ¹⁸M. Cencini, P. Muratore-Ginanneschi, and A. Vulpiani, "Nonlinear superposition of direct and inverse cascades in two-dimensional turbulence forced at large and small scales," *Phys. Rev. Lett.* **107**, 174502 (2011).
- ¹⁹M. A. Rutgers, "Forced 2D turbulence: Experimental evidence of simultaneous inverse energy and forward enstrophy cascades," *Phys. Rev. Lett.* **81**, 2244 (1998).
- ²⁰A. Belmonte, W. I. Goldburg, H. Kellay, M. A. Rutgers, B. Martin, and X. L. Wu, "Velocity fluctuations in a turbulent soap film: The third moment in two dimensions," *Phys. Fluids* **11**, 1196 (1999).
- ²¹M. Rivera, P. Vorobieff, and R. E. Ecke, "Turbulence in flowing soap films: Velocity, vorticity, and thickness fields," *Phys. Rev. Lett.* **81**, 1417 (1998).
- ²²M. Rivera and X.-L. Wu, "External dissipation in driven two-dimensional turbulence," *Phys. Rev. Lett.* **85**, 976 (2000).
- ²³M. Rivera and X.-L. Wu, "Homogeneity and the inertial range in driven two-dimensional turbulence," *Phys. Fluids* **14**, 3098 (2002).
- ²⁴C. H. Bruneau and H. Kellay, "Experiments and direct numerical simulations of two-dimensional turbulence," *Phys. Rev. E* **71**, 046305 (2005).
- ²⁵M. K. Rivera, H. Aluie, and R. E. Ecke, "The direct enstrophy cascade of two-dimensional soap film flows," *Phys. Fluids* **26**, 055105 (2014).
- ²⁶J. Paret and P. Tabeling, "Experimental observation of the two-dimensional inverse energy cascade," *Phys. Rev. Lett.* **79**, 4162 (1997).
- ²⁷J. Paret, M.-C. Jullien, and P. Tabeling, "Vorticity statistics in the two-dimensional enstrophy cascade," *Phys. Rev. Lett.* **83**, 3418 (1999).
- ²⁸H. Kellay and W. I. Goldburg, "Two-dimensional turbulence: A review of some recent experiments," *Rep. Prog. Phys.* **65**, 845 (2002).

- ²⁹L. M. Smith, J. R. Chasnov, and F. Waleffe, "Crossover from two- to three-dimensional turbulence," *Phys. Rev. Lett.* **77**, 2467 (1996).
- ³⁰A. Celani, S. Musacchio, and D. Vincenzi, "Turbulence in more than two and less than three dimensions," *Phys. Rev. Lett.* **104**, 184506 (2010).
- ³¹H. Xia, D. Byrne, G. Falkovich, and M. Shats, "Upscale energy transfer in thick turbulent fluid layers," *Nat. Phys.* **7**, 321 (2011).
- ³²D. Byrne, H. Xia, and M. Shats, "Robust inverse energy cascade and turbulence structure in three-dimensional layers of fluid," *Phys. Fluids* **23**, 095109 (2011).
- ³³G. Boffetta, F. De Lillo, and S. Musacchio, "Shell model for quasi-two-dimensional turbulence," *Phys. Rev. E* **83**, 066302 (2011).
- ³⁴H. Tennekes and J. L. Lumley, *A First Course in Turbulence* (MIT Press, 1972).
- ³⁵M. Chertkov, C. Connaughton, I. Kolokolov, and V. Lebedev, "Dynamics of energy condensation in two-dimensional turbulence," *Phys. Rev. Lett.* **99**, 084501 (2007).
- ³⁶H. Xia, M. Shats, and G. Falkovich, "Spectrally condensed turbulence in thin layers," *Phys. Fluids* **21**, 125101 (2009).
- ³⁷J. Laurie, G. Boffetta, G. Falkovich, I. Kolokolov, and V. Lebedev, "Universal profile of the vortex condensate in two-dimensional turbulence," *Phys. Rev. Lett.* **113**, 254503 (2014).
- ³⁸G. D. Nastrom and K. S. Gage, "A climatology of atmospheric wavenumber spectra of wind and temperature observed by commercial aircraft," *J. Atmos. Sci.* **42**, 950 (1985).
- ³⁹D. K. Lilly, "Stratified turbulence and the mesoscale variability of the atmosphere," *J. Atmos. Sci.* **40**, 749 (1983).
- ⁴⁰A. Vallgren, E. Deusebio, and E. Lindborg, "Possible explanation of the atmospheric kinetic and potential energy spectra," *Phys. Rev. Lett.* **107**, 268501 (2011).
- ⁴¹A. Campagne, B. Gallet, F. Moisy, and P.-P. Cortet, "Direct and inverse energy cascades in a forced rotating turbulence experiment," *Phys. Fluids* **26**, 125112 (2014).
- ⁴²H. K. Moffatt, "Note on the triad interactions of homogeneous turbulence," *J. Fluid Mech.* **741**, R3 (2014).
- ⁴³L. Biferale, M. Buzicotti, and M. Linkmann, e-print [arXiv:1706.02371](https://arxiv.org/abs/1706.02371) [physics.flu-dyn] (2017).
- ⁴⁴E. Deusebio, G. Boffetta, E. Lindborg, and S. Musacchio, "Dimensional transition in rotating turbulence," *Phys. Rev. E* **90**, 023005 (2014).
- ⁴⁵A. Sozza, G. Boffetta, P. Muratore-Ginanneschi, and S. Musacchio, "Dimensional transition of energy cascades in stably stratified forced thin fluid layers," *Phys. Fluids* **27**, 035112 (2015).