

The GSI Time Anomaly: Facts and Fiction

Carlo Giunti

INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino

<mailto://giunti@to.infn.it>

Neutrino Unbound: <http://www.nu.to.infn.it>


IFIC, Valencia, 3 December 2008

Outline

- The GSI Anomaly
- Neutrino Mixing?
- Interference: Double-Slit Analogy
- arXiv:0801.1465 and arXiv:0805.0435
- arXiv:0801.2121 – arXiv:0801.3262
- Quantum Beats?
- Towards the Epilogue?
- Conclusions

The GSI Experiment

Schematic layout of the secondary nuclear beam facility at GSI




[Litvinov et al, nucl-ex/0509019]

SIS: Heavy Ion Synchrotron

FRS: FRagment Separator

ESR: Experiment Storage Ring




Schottky Mass Spectrometry

- ▶ Stored ions circulate in ESR with revolution frequencies ~ 2 MHz
- ▶ At each turn they induce mirror charges on two electrodes
- ▶ Revolution frequency spectra provide information about q/m :



$$f = \frac{\omega}{2\pi} = \frac{qB}{2\pi m\gamma}$$

- ▶ Area of each frequency peak is proportional to number of stored ions




[Litvinov et al, nucl-ex/0509019]

Praseodymium


atomic number	59	140.908	atomic weight
symbol	Pr		acid-base properties of higher-valence oxides
electron configuration	[Xe]4f ³ 6s ²		crystal structure
name	praseodymium		physical state at 20° C (68° F)

	weakly basic		solid
	hexagonal		rare earth elements lanthanide elements



© 2006 Encyclopædia Britannica, Inc.



[Litvinov et al, PLB 664 (2008) 168]





Cerium



atomic number	58	140.116	atomic weight
symbol	Ce		acid-base properties of higher-valence oxides
electron configuration	[Xe]4f ² 6s ²		crystal structure
name	cerium		physical state at 20° C (68° F)

	weakly basic		solid
	cubic, face centred		rare earth elements lanthanide elements

© 2006 Encyclopædia Britannica, Inc.


Promethium

atomic number	61	(145)	atomic weight
symbol	Pm		acid-base properties of higher-valence oxides
electron configuration	[Xe]4f ⁵ 6s ²		crystal structure
name	promethium		physical state at 20° C (68° F)

	weakly basic		synthetically prepared
	hexagonal		rare earth elements lanthanide elements



() indicates the mass of the longest-lived isotope

© 2006 Encyclopædia Britannica, Inc.




[Litvinov et al, PLB 664 (2008) 168]

Neodymium




atomic number	60	144.24	atomic weight
symbol	Nd		acid-base properties of higher-valence oxides
electron configuration	[Xe]4f ⁴ 6s ²		crystal structure
name	neodymium		physical state at 20° C (68° F)

	weakly basic		solid
	hexagonal		rare earth elements lanthanide elements


© 2006 Encyclopædia Britannica, Inc.




- ▶ About six initial $^{140}_{59}\text{Pr}^{58+}$ ions ($f = qB/2\pi m\gamma$)
- ▶ Two decayed via nuclear electron capture into $^{140}_{58}\text{Ce}^{58+}$
- ▶ Seen because $\Delta q = 0 \Rightarrow \Delta f/f = -\Delta m/m$ (small)
- ▶ Other decayed via β^+ decay ($\Delta q = -1 \Rightarrow \Delta f \sim -150$ kHz) or were lost (interactions with residual gas)




[Litvinov et al, PLB 664 (2008) 168]



[Litvinov et al, PLB 664 (2008) 168]



[Litvinov et al, PLB 664 (2008) 168]



[Litvinov et al, PLB 664 (2008) 168]

$$(1) \quad \frac{dN_{EC}(t)}{dt} = \lambda_{EC} N(t) = \lambda_{EC} N(0) e^{-\lambda t}$$

$$(2) \quad \frac{dN_{EC}(t)}{dt} = \tilde{\lambda}_{EC}(t) N(t) = \tilde{\lambda}_{EC}(t) N(0) e^{-\lambda t}$$

$$\lambda = \lambda_{EC} + \lambda_{\beta^+} + \lambda_{loss} \quad \tilde{\lambda}_{EC}(t) = \lambda_{EC} [1 + a \cos(\omega t + \phi)]$$

Fit parameters of $^{140}_{59}\text{Pr}$ data					
Eq.	$N_0 \lambda_{EC}$	λ	a	ω	χ^2/DoF
(1)	34.9(18)	0.00138(10)	-	-	107.2/73
(2)	35.4(18)	0.00147(10)	0.18(3)	0.89(1)	67.18/70
Fit parameters of $^{142}_{61}\text{Pm}$ data					
Eq.	$N_0 \lambda_{EC}$	λ	a	ω	χ^2/DoF
(1)	46.8(40)	0.0240(42)	-	-	63.77/38
(2)	46.0(39)	0.0224(41)	0.23(4)	0.89(3)	31.82/35

$$T(^{140}_{59}\text{Pr}^{58+}) = 7.06 \pm 0.08 \text{ s} \quad T(^{142}_{61}\text{Pm}^{60+}) = 7.10 \pm 0.22 \text{ s}$$

$$\langle a \rangle = 0.20 \pm 0.02$$

[Litvinov et al, PLB 664 (2008) 168]

Neutrino Mixing?

[Litvinov et al, PLB 664 (2008) 168]

$$l_i \rightarrow l_f + \nu_e \quad \nu_e = \cos \vartheta_{\text{SOL}} \nu_1 + \sin \vartheta_{\text{SOL}} \nu_2$$

PROPOSED EXPLANATION: INTERFERENCE OF ν_1 AND ν_2

Initial Ion: Momentum $\vec{P} = 0$, Energy E

Massive ν_k : Momentum \vec{p}_k , Energy $E_k = \sqrt{p_k^2 + m_k^2}$

Final Ion: Momentum $-\vec{p}_k$, Energy $M + p_k^2/2M$

$$E_1 + M + p_1^2/2M = E \quad E_2 + M + p_2^2/2M = E$$

$$\Delta E \equiv E_2 - E_1 \simeq \frac{\Delta m^2}{2M} \quad \Delta m^2 \equiv m_2^2 - m_1^2$$

massive neutrino energy difference: $\Delta E \equiv E_2 - E_1 \simeq \frac{\Delta m^2}{2M}$

$$\Delta m^2 = \Delta m_{\text{SOL}}^2 \simeq 8 \times 10^{-5} \text{ eV}^2 \quad M \simeq 140 \text{ amu} \simeq 130 \text{ GeV}$$


$$\Delta E \simeq 3.1 \times 10^{-16} \text{ eV}$$

$$T = \frac{2\pi}{\Delta E} \gamma \simeq 19.1 \text{ s} \quad \gamma = 1.43$$

about 3 times larger than $T_{\text{GSI}} \simeq 7 \text{ s}$

CAN INTERFERENCE IN FINAL STATE AFFECT DECAY RATE?

Interference: Double-Slit Analogy



- ▶ Decay rate of \mathbb{I} corresponds to fraction of intensity of incoming wave which crosses the barrier
- ▶ Fraction of intensity of the incoming wave which crosses the barrier depends on the sizes of the holes
- ▶ It does not depend on interference effects which occur after the wave has passed through the barrier
- ▶ Analogy: decay rate of \mathbb{I} cannot depend on interference of ν_1 and ν_2 which occurs after decay has happened \iff CAUSALITY!

Causality

INTERFERENCE OF
COHERENT ENERGY STATES

(ν_1 AND ν_2)

OCCURRING **AFTER** THE DECAY

(flavor neutrino oscillations)

CANNOT AFFECT THE DECAY RATE

Cross Sections and Decay Rates are always summed incoherently over different final channels:

$$\mathbb{I} \rightarrow \mathbb{F}_1, \quad \mathbb{I} \rightarrow \mathbb{F}_2, \quad \dots \quad \Longrightarrow \quad P_{\mathbb{I} \rightarrow \mathbb{F}} = \sum_k P_{\mathbb{I} \rightarrow \mathbb{F}_k}$$

entangled final state: $|\mathbb{F}\rangle = \sum_k A_k |\mathbb{F}_k\rangle$

$$|\mathbb{F}\rangle \propto (\mathbf{S} - \mathbf{1}) |\mathbb{I}\rangle \quad \Longrightarrow \quad A_k \propto \langle \mathbb{F}_k | (\mathbf{S} - \mathbf{1}) |\mathbb{I}\rangle = \langle \mathbb{F}_k | \mathbf{S} | \mathbb{I}\rangle$$

$$\langle \mathbb{F} | \mathbb{F} \rangle = 1 \quad \Longleftrightarrow \quad A_k = \frac{\langle \mathbb{F}_k | \mathbf{S} | \mathbb{I}\rangle}{\left(\sum_j |\langle \mathbb{F}_j | \mathbf{S} | \mathbb{I}\rangle|^2 \right)^{1/2}}$$

$$\begin{aligned} P_{\mathbb{I} \rightarrow \mathbb{F}} &= |\langle \mathbb{F} | \mathbf{S} | \mathbb{I}\rangle|^2 = \left| \sum_k A_k^* \langle \mathbb{F}_k | \mathbf{S} | \mathbb{I}\rangle \right|^2 = \left(\frac{\sum_k |\langle \mathbb{F}_k | \mathbf{S} | \mathbb{I}\rangle|^2}{\left(\sum_j |\langle \mathbb{F}_j | \mathbf{S} | \mathbb{I}\rangle|^2 \right)^{1/2}} \right)^2 \\ &= \sum_k |\langle \mathbb{F}_k | \mathbf{S} | \mathbb{I}\rangle|^2 = \sum_k P_{\mathbb{I} \rightarrow \mathbb{F}_k} \end{aligned}$$


coherent character of final state is irrelevant for interaction probability!

arXiv:0801.1465 and arXiv:0805.0435

H.J. Lipkin

- ▶ Causality is violated explicitly
- ▶ arXiv:0801.1465: The difference in momentum δp_ν between the two neutrino eigenstates with the same energy produces a small initial momentum change $\delta P \dots$
- ▶ arXiv:0805.0435: Since the time dependence depends only on the propagation of the initial state, it is independent of the final state, which is created only at the decay point. Thus there is no violation of causality.
- ▶ But in calculation of effect: The phase difference at a time t between states produced by the neutrino mass difference on the motion of the initial ion in the laboratory frame with velocity $V = (P/E)$ is

$$\delta\phi \approx -\delta E \cdot t = \Delta m^2/2E$$



- ▶ $\mathbb{I} \rightarrow \mathbb{F} + \nu$ decay rate in time-dependent perturbation theory

with final neutrino state $|\nu\rangle = \sum_k |\nu_k\rangle$

- ▶ Not even properly normalized to describe one particle:

$$\langle \nu_j | \nu_k \rangle = \delta_{jk} \implies \langle \nu | \nu \rangle = 3$$

- ▶ Different from standard electron neutrino state

$$|\nu_e\rangle = \sum_k U_{ek}^* |\nu_k\rangle$$

Time-Dependent Perturbation Theory

$$P_{\mathbb{I} \rightarrow \mathbb{F} + \nu}(t) = \left| \int_0^t d\tau \langle \nu, \mathbb{F} | \mathcal{H}_W(\tau) | \mathbb{I} \rangle \right|^2 = \left| \sum_k \int_0^t d\tau \langle \nu_k, \mathbb{F} | \mathcal{H}_W(\tau) | \mathbb{I} \rangle \right|^2$$

$$\mathcal{H}_W(t) = \int d^3x \mathcal{H}_W(x)$$

Effective Four-Fermion Interaction Hamiltonian

$$\begin{aligned} \mathcal{H}_W(x) &= \frac{G_F}{\sqrt{2}} \cos \theta_C \bar{\nu}_e(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) \\ &= \frac{G_F}{\sqrt{2}} \cos \theta_C \sum_k U_{ek}^* \bar{\nu}_k(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) \end{aligned}$$

$$\langle \nu_k, \mathbb{F} | \mathcal{H}_W(\tau) | \mathbb{I} \rangle = U_{ek}^* e^{i\Delta E_k \tau} T_k \quad \text{with} \quad \Delta E_k = E_k + E_{\mathbb{F}} - E_{\mathbb{I}}$$

$$\int_0^t d\tau e^{i\Delta E_k \tau} = e^{i\Delta E_k t/2} \frac{\sin(\Delta E_k t/2)}{\Delta E_k/2} \xrightarrow{\Delta E_k t \gg 1} 2\pi \delta(\Delta E_k) e^{i\Delta E_k t/2}$$

$$P_{\mathbb{I} \rightarrow \mathbb{F} + \nu}(t) = 4\pi^2 \left| \sum_k U_{ek}^* e^{i\Delta E_k t} \delta(\Delta E_k) T_k \right|^2$$

$$T_k \simeq T_j$$

$\delta(\Delta E_k)$ satisfied by wave packet

$$P_{\mathbb{I} \rightarrow \mathbb{F} + \nu}(t) \propto \left| \sum_k U_{ek}^* e^{i\Delta E_k t} \right|^2$$

Two-Neutrino Mixing

$$\begin{aligned} P_{\mathbb{I} \rightarrow \mathbb{F} + \nu}(t) &\propto \left| \cos \vartheta e^{i\Delta E_1 t} + \sin \vartheta e^{i\Delta E_2 t} \right|^2 = 1 + \sin 2\vartheta \cos\left(\frac{\Delta E t}{2}\right) \\ &= 1 + \sin 2\vartheta \cos\left(\frac{\Delta m^2 t}{4M}\right) \end{aligned}$$

$$\Delta E = \Delta E_2 - \Delta E_1 = E_2 - E_1 = \frac{\Delta m^2}{2M}$$

▶ Standard QFT: $P_{\mathbb{I} \rightarrow \mathbb{F} + \nu} = |\langle \nu, \mathbb{F} | S | \mathbb{I} \rangle|^2 = \left| \sum_k \langle \nu_k, \mathbb{F} | S | \mathbb{I} \rangle \right|^2$

▶ S-matrix operator at first order in perturbation theory:

$$S = 1 - i \int d^4x \mathcal{H}_W(x)$$

▶ Effective four-fermion interaction Hamiltonian:

$$\begin{aligned} \mathcal{H}_W(x) &= \frac{G_F}{\sqrt{2}} \cos \theta_C \bar{\nu}_e(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) \\ &= \frac{G_F}{\sqrt{2}} \cos \theta_C \sum_k U_{ek}^* \bar{\nu}_k(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) \end{aligned}$$

▶ $\langle \nu_k, \mathbb{F} | S | \mathbb{I} \rangle = U_{ek}^* \mathcal{M}_k$ with

$$\mathcal{M}_k = -i \frac{G_F}{\sqrt{2}} \cos \theta_C \int d^4x \langle \nu_k, \mathbb{F} | \bar{\nu}_k(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) | \mathbb{I} \rangle$$

▶ $P_{\mathbb{I} \rightarrow \mathbb{F} + \nu} = \left| \sum_k U_{ek}^* \mathcal{M}_k \right|^2$ different from standard $P = \sum_k |U_{ek}|^2 |\mathcal{M}_k|^2$

- **Check:** in the limit of massless neutrinos decay probability should reduce to the Standard Model decay probability

$$P_{\text{SM}} = |\mathcal{M}_{\text{SM}}|^2$$

with

$$\mathcal{M}_{\text{SM}} = -i \frac{G_F}{\sqrt{2}} \cos \theta_C \int d^4x \langle \nu_e, \mathbb{F} | \bar{\nu}_e(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) | \mathbb{I} \rangle$$

where ν_e is the Standard Model massless electron neutrino

$$\mathcal{M}_k = -i \frac{G_F}{\sqrt{2}} \cos \theta_C \int d^4x \langle \nu_k, \mathbb{F} | \bar{\nu}_k(x) \gamma_\rho (1 - \gamma^5) e(x) \bar{n}(x) \gamma^\rho (1 - g_A \gamma^5) p(x) | \mathbb{I} \rangle$$

$$\mathcal{M}_k \xrightarrow{m_k \rightarrow 0} \mathcal{M}_{\text{SM}}$$

$$P_{\mathbb{I} \rightarrow \mathbb{F} + \nu} = \left| \sum_k U_{ek}^* \mathcal{M}_k \right|^2 \xrightarrow{m_k \rightarrow 0} |\mathcal{M}_{\text{SM}}|^2 \left| \sum_k U_{ek}^* \right|^2 \neq P_{\text{SM}}$$

WRONG!

- ▶ Correct normalized final neutrino state ($\langle \nu_e | \nu_e \rangle = 1$):

$$\begin{aligned}
 |\nu_e\rangle &= \left(\sum_j |\langle \nu_j, \mathbb{F} | S | \mathbb{I} \rangle|^2 \right)^{-1/2} \sum_k |\nu_k\rangle \langle \nu_k, \mathbb{F} | S | \mathbb{I} \rangle \\
 &= \left(\sum_j |U_{ej}|^2 |\mathcal{M}_j|^2 \right)^{-1/2} \sum_k U_{ek}^* \mathcal{M}_k |\nu_k\rangle
 \end{aligned}$$

- ▶ Standard decay probability:

$$P_{\mathbb{I} \rightarrow \mathbb{F} + \nu_e} = |\langle \nu_e, \mathbb{F} | S | \mathbb{I} \rangle|^2 = \sum_k |\langle \nu_k, \mathbb{F} | S | \mathbb{I} \rangle|^2 = \sum_k |U_{ek}|^2 |\mathcal{M}_k|^2$$

$$P_{\mathbb{I} \rightarrow \mathbb{F} + \nu_e} \xrightarrow{m_k \rightarrow 0} P_{\text{SM}}$$

- ▶ In experiments which are not sensitive to the differences of the neutrino masses, as neutrino oscillation experiments,

$$\mathcal{M}_k \simeq \overline{\mathcal{M}} \quad \Longrightarrow \quad |\nu_e\rangle = \sum_k U_{ek}^* |\nu_k\rangle$$

Time-Dependent Perturbation Theory?

not appropriate because electron capture and decay are interrupted by

Schottky Mass Spectrometry

with ESR revolution frequency ~ 2 MHz, i.e. every

$$\sim 5 \times 10^{-7} \text{ s}$$

much smaller than ion lifetime

$$T_{1/2}(^{140}_{59}\text{Pr}) \simeq 3.39 \text{ m}$$

$$T_{1/2}(^{142}_{61}\text{Pm}) \simeq 40.5 \text{ s}$$

and period of anomalous oscillations $T \simeq 7 \text{ s}$

interaction time:
$$t_W \sim \frac{\hbar}{m_W} \simeq \frac{6.6 \times 10^{-22} \text{ MeV s}}{8.0 \times 10^4 \text{ MeV}} \sim 10^{-26} \text{ s}$$


$t \gg t_W$ in Time-Dependent Perturbation Theory



Quantum Field Theory result

Quantum Beats?

- ▶ GSI time anomaly can be due to interference effects in **initial** state
- ▶ Two coherent energy states of the decaying ion \implies **Quantum Beats**



$$\mathbb{I} = \mathcal{A}_1 \mathbb{I}_1 + \mathcal{A}_2 \mathbb{I}_2$$

INTERFERENCE = QUANTUM BEATS

INTERFERENCE = OSCILLATIONS

$$\nu_e = \cos \vartheta \nu_1 + \sin \vartheta \nu_2$$

A vector diagram showing a vertical purple vector labeled \mathbb{I} at the top and \mathbb{F} at the bottom. A red vector branches off from the purple vector, pointing downwards and to the right. The angle between the purple vector and the red vector is labeled ϑ .

- ▶ Incoming waves interfere at holes in barrier
- ▶ **Causality**: interference due to different phases of incoming waves developed during propagation **before** reaching the barrier

Causality

INTERFERENCE OF
COHERENT ENERGY STATES
OCCURRING **BEFORE** THE DECAY
CAN AFFECT THE DECAY RATE

Quantum Beats




Fig. 1 Diagram of the four level system. A photon is absorbed by the ground state $|g\rangle$ and excites a superposition of states $|a\rangle$ and $|b\rangle$ whose energy separation is $\Delta E = \hbar\omega_{ab}$. Emission of a second photon leaves the system in the final state $|f\rangle$.

$$I_{\text{fl}}(t) \propto |\mu_{ag}|^2 |\mu_{fa}|^2 e^{-\gamma_a t} + |\mu_{bg}|^2 |\mu_{fb}|^2 e^{-\gamma_b t} + |\mu_{ag}\mu_{bg}\mu_{fa}\mu_{fb}| e^{-(\gamma_a + \gamma_b)t/2} \cos(\omega_{ab}t + \theta). \quad (4)$$

Examination of this expression shows that it consists of two parts, one incoherent term (first two terms) describing the independent decays of the two states $|a\rangle$ and $|b\rangle$ and one coherent or cross term (last term) which decays at the average rate of the two states and, most importantly, is modulated at the angular frequency ω_{ab} . The modulation frequency is the difference of the two angular frequencies in eqn. (2), *i.e.* $\omega_{ab} = |\omega_a - \omega_b|$, and the coherent term in eqn. (4) is therefore termed the quantum beat. The angle θ is included in eqn. (4) to describe the phase of the quantum beat, which depends on a number of factors such as the excitation and detection polarisations and transitions. When the transition moments and decay rates are equal, as is often the case, a particularly simple expression is derived for the four level system. In this case eqn. (4) becomes

$$I_{\text{fl}}(t) \propto [1 + \cos(\omega_{ab}t + \theta)]e^{-\gamma t}, \quad (5)$$

clearly illustrating the contributions of the incoherent and coherent terms to the fluorescence decay. In this special case the quantum beat is 100% modulated. It is important to point out

[Carter, Huber, Quantum beat spectroscopy in chemistry, Chem. Soc. Rev., 29 (2000) 305]




Fig. 2 Zeeman quantum beat recorded for the $R(0)$ line of the $17U$ transition in CS_2 in an external field of ~ 15 Gauss. The laser polarisation was perpendicular to the magnetic field direction and prepares a coherence between the $M = \pm 1$ sublevels as shown in the level diagram. This is manifested by a single quantum beat on the fluorescence decay; the real part of the Fourier transform is also shown. The less than 100% modulation, which is observed in virtually all quantum beat measurements in molecules, is due to incoherent emission from the excited states.




Fig. 6 Nuclear hyperfine quantum beats recorded for the $P_{2v}/Q_1(3/2)$ line in a vibrational band of the $A^2\Sigma^+ - X^2\Pi$ transition in the Ar-OD van der Waals complex. The inset shows the fluorescence decay which exhibits weakly modulated quantum beats. Following Fourier transformation the beat frequencies between hyperfine levels in the $A^2\Sigma^+$ state are clearly visible.

[Carter, Huber, Quantum beat spectroscopy in chemistry, Chem. Soc. Rev., 29 (2000) 305]

- ▶ Quantum beats in GSI experiment can be due to interference of two coherent energy states of the decaying ion which develop different phases before the decay
- ▶ Coherence is preserved for a long time if measuring apparatus which monitors the ions with frequency ~ 2 MHz does not distinguish between the two states

$$\text{▶ } |\mathbb{I}(t=0)\rangle = \mathcal{A}_1 |\mathbb{I}_1\rangle + \mathcal{A}_2 |\mathbb{I}_2\rangle \quad (|\mathcal{A}_1|^2 + |\mathcal{A}_2|^2 = 1)$$

$$\Gamma = \Gamma_1 \simeq \Gamma_2 \implies |\mathbb{I}(t)\rangle = \left(\mathcal{A}_1 e^{-iE_1 t} |\mathbb{I}_1\rangle + \mathcal{A}_2 e^{-iE_2 t} |\mathbb{I}_2\rangle \right) e^{-\Gamma t/2}$$

$$P_{\text{EC}}(t) = |\langle \nu_e, \mathbb{F} | S | \mathbb{I}(t) \rangle|^2 = [1 + A \cos(\Delta E t + \varphi)] \bar{P}_{\text{EC}} e^{-\Gamma t}$$

$$A \equiv 2|\mathcal{A}_1||\mathcal{A}_2|, \quad \Delta E \equiv E_2 - E_1, \quad \bar{P}_{\text{EC}} = |\langle \nu_e, \mathbb{F} | S | \mathbb{I}_1 \rangle|^2 \simeq |\langle \nu_e, \mathbb{F} | S | \mathbb{I}_2 \rangle|^2$$

$$\frac{dN_{\text{EC}}(t)}{dt} = N(0) [1 + A \cos(\Delta E t + \varphi)] \bar{\Gamma}_{\text{EC}} e^{-\Gamma t}$$

$$\frac{dN_{\text{EC}}(t)}{dt} = N(0) [1 + A \cos(\Delta E t + \varphi)] \bar{\Gamma}_{\text{EC}} e^{-\Gamma t}$$

$$\Delta E({}_{59}^{140}\text{Pr}^{58+}) = (5.86 \pm 0.07) \times 10^{-16} \text{ eV}, \quad A({}_{59}^{140}\text{Pr}^{58+}) = 0.18 \pm 0.03$$


$$\Delta E({}_{61}^{142}\text{Pm}^{60+}) = (5.82 \pm 0.18) \times 10^{-16} \text{ eV}, \quad A({}_{61}^{142}\text{Pm}^{60+}) = 0.23 \pm 0.04$$

$$A \equiv 2|\mathcal{A}_1||\mathcal{A}_2|$$

- ▶ Energy splitting is extremely small
- ▶ $|\mathcal{A}_1|^2/|\mathcal{A}_2|^2 \sim 1/99$ or $|\mathcal{A}_2|^2/|\mathcal{A}_1|^2 \sim 1/99$
- ▶ It is difficult to find an appropriate mechanism

Hyperfine Splitting

smallest known energy splitting



[Litvinov et al, PRL 99 (2007) 262501]

$$\Delta E \sim 10^{-6} \text{ eV} \quad \Longrightarrow \quad T \sim 10^{-9} \text{ s} \quad f \sim \text{GHz}$$

too large to explain the GSI anomaly

$$T_{\text{GSI}} \simeq 7 \text{ s} \quad f_{\text{GSI}} \simeq 0.14 \text{ Hz} \quad \Delta E_{\text{GSI}} = 2\pi/T_{\text{GSI}} \simeq 6 \times 10^{-16} \text{ eV}$$

feeling of smallness of $\Delta E_{\text{GSI}} \simeq 6 \times 10^{-16} \text{ eV}$

$$\mu_{\text{N}} B_{\oplus} \simeq \left(3 \times 10^{-12} \text{ eV G}^{-1} \right) (0.5 \text{ G}) = 1.5 \times 10^{-12} \text{ eV}$$

$$\Delta E_{\text{GSI}} \simeq 4 \times 10^{-4} \mu_{\text{N}} B_{\oplus}$$

Towards the Epilogue?

Berkeley Experiment – arXiv:0807.0649

- ▶ EC: $^{142}\text{Pm} \rightarrow ^{142}\text{Nd} + \nu_e$ ^{142}Pm in an aluminum foil
no oscillations at a level 31 times smaller than GSI
- ▶ Reanalysis of old $^{142}\text{Eu} \rightarrow ^{142}\text{Sm} + \nu_e$ EC data \implies no oscillations
- ▶ Differences with GSI Experiment:
 - ▶ neutral atoms
 - ▶ stopped atoms

- ▶ “orbital electron-capture decays of neutral ^{142}Pm atoms implanted into the lattice of a solid do not fulfil the constraints of true two-body beta decays, since momentum as well as energy of the final state are distributed among three objects, namely the electron neutrino, the recoiling daughter atom and the lattice phonons.”
- ▶ $\Delta E \simeq 8 \cdot 10^{-16} \text{ eV} \ll 10^{-3} \text{ eV}$ typical phonon energy.
“the modulations could be washed out”
- ▶ “A measurement of the EC-decay of helium-like ^{142}Pm ions should reveal the (probably small) differences to the EC-decay of hydrogen-like ions (such time-resolved measurements are planned, but not yet performed). And, without doubt, the outcome of the three-body β^+ decay of ^{142}Pm is crucial for the interpretation of the GSI data. The-not simple-evaluation of this data is still in progress.”

! remarks based on wrong neutrino interference hypothesis !

Munich Group + F. Bosch (GSI) – arXiv:0807.3297

- ▶ $^{180}\text{Re} \rightarrow ^{180}\text{W} + \nu_e$ ^{180}Re in a tantalum foil
no oscillations at a level more than 10 times smaller than GSI
 - ▶ “The GSI oscillations are not observable in a conventional experiment with radioactive atoms in a solid environment but must have to do with the **unique conditions in the GSI experiment** where hydrogen-like ions are moving independently in a storage ring and decaying directly by a true two-body decay to a long-lived (ground-) state.”
 - ▶ “The recoiling daughter atom has **not a well defined momentum**, because it moves in a lattice and **can only assume momenta allowed by the phonon spectrum**. Therefore the effects causing oscillations in the EC decay to bare ions coasting in a storage ring could be washed out in a standard experiment.”
- ! vague statements based on wrong neutrino interference hypothesis !

Spin-rotation coupling in non-exponential decay of hydrogenlike heavy ions

14 November 2008

We discuss a model in which a recently reported modulation in the decay of the hydrogenlike ions $^{140}\text{Pr}^{58+}$ and $^{142}\text{Pm}^{60+}$ arises from the coupling of rotation to the spin of electron and nuclei (Thomas precession). A similar model describes the electron modulation in muon $g - 2$ experiments correctly. Agreement with the GSI experimental results is obtained for the current QED-values of the bound electron g -factors, $g(^{140}\text{Pr}^{58+}) = 1.872$ and $g(^{142}\text{Pm}^{60+}) = 1.864$, if the Lorentz factor of the bound electron is ~ 1.88 . The latter is fixed by either of the two sets of experimental data. The model predicts that the modulation is not observable if the motion of the ions is linear, or if the ions are stopped in a target.

Conclusions

- ▶ **Interference:** due to phase difference of two incoming waves
- ▶ **Causality:** there cannot be interference of waves before they exist
- ▶ The GSI ion lifetime anomaly **cannot** be due to interference of decay product before the decay product start to exist (neutrino mixing in the final state)
- ▶ The GSI ion lifetime anomaly **can** be due to interference of two energy states of the decaying ion: **Quantum Beats**
- ▶ No known mechanism, because
 - ▶ Energy splitting of the two energy states: $\Delta E \sim 6 \times 10^{-16} \text{ eV}$
 - ▶ Ratio of probabilities of the two energy states: 1/99