

Status of eV Sterile Neutrinos

Carlo Giunti

INFN, Torino, Italy

PPNS 2018

International Workshop on Particle Physics at Neutron Sources

24-26 May 2018, ILL, Grenoble, France


Neutrino Oscillations

$$\nu_{\alpha L} = \sum_k U_{\alpha k} \nu_{kL} \quad \alpha = e, \mu, \tau$$



$$P_{\nu_\alpha \rightarrow \nu_\beta}(L, E) = \delta_{\alpha\beta} - 4 \underbrace{\sum_{k>j} \text{Re} [U_{\alpha k}^* U_{\beta k} U_{\alpha j} U_{\beta j}^*] \sin^2 \left(\frac{\Delta m_{kj}^2 L}{4E} \right)}_{\text{CP conserving}} \\ + 2 \underbrace{\sum_{k>j} \text{Im} [U_{\alpha k}^* U_{\beta k} U_{\alpha j} U_{\beta j}^*] \sin \left(\frac{\Delta m_{kj}^2 L}{2E} \right)}_{\text{CP violating}}$$

- ▶ Squared-mass differences: $\Delta m_{kj}^2 = m_k^2 - m_j^2$
- ▶ Mixing: $U_{\alpha k}^* U_{\beta k} U_{\alpha j} U_{\beta j}^*$ quartic rephasing invariants
- ▶ Jarlskog invariant: $J_{\text{CP}} = \text{Im} [U_{\alpha k}^* U_{\beta k} U_{\alpha j} U_{\beta j}^*]$

Three-Neutrino Mixing Paradigm



Beyond Three-Neutrino Mixing: Sterile Neutrinos



$$N_{\nu_{\text{active}}}^{\text{LEP}} = 2.9840 \pm 0.0082$$

Terminology: a eV-scale sterile neutrino
means: a eV-scale massive neutrino which is mainly sterile

Sterile Neutrinos from Physics Beyond the SM

- ▶ Neutrinos are special in the Standard Model: the only **neutral fermions**
- ▶ Active left-handed neutrinos can mix with non-SM singlet fermions often called **right-handed neutrinos**
- ▶ Light left-handed anti- ν_R are **light sterile neutrinos**

$$\nu_R^c \rightarrow \nu_{sL} \quad (\text{left-handed})$$

- ▶ Sterile means **no standard model interactions**

[Pontecorvo, Sov. Phys. JETP 26 (1968) 984]

- ▶ Active neutrinos (ν_e, ν_μ, ν_τ) can oscillate into light sterile neutrinos (ν_s)
- ▶ Observables:
 - ▶ Disappearance of active neutrinos (neutral current deficit) \leftarrow CE ν NS
 - ▶ Indirect evidence through combined fit of data (current indication)
- ▶ Short-baseline anomalies + 3ν -mixing:

$$\Delta m_{21}^2 \ll |\Delta m_{31}^2| \ll |\Delta m_{41}^2| \leq \dots$$

ν_1	ν_2	ν_3	ν_4	\dots
ν_e	ν_μ	ν_τ	ν_{s1}	\dots

Effective 3+1 SBL Oscillation Probabilities

Appearance ($\alpha \neq \beta$)

$$P_{\nu_\alpha \rightarrow \nu_\beta}^{\text{SBL}} \simeq \sin^2 2\vartheta_{\alpha\beta} \sin^2 \left(\frac{\Delta m_{41}^2 L}{4E} \right)$$

$$\sin^2 2\vartheta_{\alpha\beta} = 4|U_{\alpha 4}|^2 |U_{\beta 4}|^2$$

$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} & U_{\mu 4} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} & U_{\tau 4} \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} \end{pmatrix}_{\text{SBL}}$$

- ▶ 6 mixing angles
- ▶ 3 Dirac CP phases
- ▶ 3 Majorana CP phases

Disappearance

$$P_{\nu_\alpha \rightarrow \nu_\alpha}^{\text{SBL}} \simeq 1 - \sin^2 2\vartheta_{\alpha\alpha} \sin^2 \left(\frac{\Delta m_{41}^2 L}{4E} \right)$$

$$\sin^2 2\vartheta_{\alpha\alpha} = 4|U_{\alpha 4}|^2 (1 - |U_{\alpha 4}|^2)$$

- ▶ CP violation is not observable in SBL experiments!
- ▶ Observable in LBL accelerator exp. sensitive to Δm_{ATM}^2 [de Gouvea et al, PRD 91 (2015) 053005, PRD 92 (2015) 073012, arXiv:1605.09376; Palazzo et al, PRD 91 (2015) 073017, PLB 757 (2016) 142; Kayser et al, JHEP 1511 (2015) 039, JHEP 1611 (2016) 122] and solar exp. sensitive to Δm_{SOL}^2 [Long, Li, CG, PRD 87, 113004 (2013) 113004]

3+1: Appearance vs Disappearance

- Amplitude of ν_e disappearance:

$$\sin^2 2\vartheta_{ee} = 4|U_{e4}|^2 (1 - |U_{e4}|^2) \simeq 4|U_{e4}|^2$$

- Amplitude of ν_μ disappearance:

$$\sin^2 2\vartheta_{\mu\mu} = 4|U_{\mu 4}|^2 (1 - |U_{\mu 4}|^2) \simeq 4|U_{\mu 4}|^2$$

- Amplitude of $\nu_\mu \rightarrow \nu_e$ transitions:

$$\sin^2 2\vartheta_{e\mu} = 4|U_{e4}|^2 |U_{\mu 4}|^2 \simeq \frac{1}{4} \sin^2 2\vartheta_{ee} \sin^2 2\vartheta_{\mu\mu}$$

quadratically suppressed for small $|U_{e4}|^2$ and $|U_{\mu 4}|^2$



Appearance-Disappearance Tension


[Okada, Yasuda, IJMPA 12 (1997) 3669; Bilenky, CG, Grimus, EPJC 1 (1998) 247]

Gallium Anomaly

Gallium Radioactive Source Experiments: GALLEX and SAGE




Test of Solar ν_e Detection:



$$\langle L \rangle_{\text{GALLEX}} = 1.9 \text{ m} \quad \langle L \rangle_{\text{SAGE}} = 0.6 \text{ m}$$

$$\Delta m^2_{\text{SBL}} \gtrsim 1 \text{ eV}^2 \gg \Delta m^2_{\text{ATM}}$$

► ${}^3\text{He} + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + {}^3\text{H}$ cross section measurement



$\approx 2.9\sigma$ deficit


[SAGE, PRC 73 (2006) 045805; PRC 80 (2009) 015807;
Laveder et al, Nucl.Phys.Proc.Suppl. 168 (2007) 344,
MPLA 22 (2007) 2499, PRD 78 (2008) 073009,
PRC 83 (2011) 065504]


Reactor Electron Antineutrino Anomaly

[Mention et al, PRD 83 (2011) 073006]

New reactor $\bar{\nu}_e$ fluxes

[Mueller et al, PRC 83 (2011) 054615; Huber, PRC 84 (2011) 024617]








$$\Delta m^2_{SBL} \gtrsim 0.5 \text{ eV}^2 \gg \Delta m^2_{\text{ATM}}$$

- SBL oscillations are averaged at the Daya Bay, RENO, and Double Chooz near detectors \Rightarrow no spectral distortion


Reactor Antineutrino 5 MeV Bump



- ▶ Cannot be explained by neutrino oscillations (SBL oscillations are averaged in RENO, DC, DB).
- ▶ It is likely due to a theoretical miscalculation of the spectrum. (Or to detector energy nonlinearity [Mention et al, PLB 773 (2017) 307]).
- ▶ $\sim 3\%$ effect on total flux, but if it is an excess it increases the anomaly!
- ▶ No post-bump complete calculation of the neutrino fluxes.
- ▶ Nominal Saclay-Huber flux calculation uncertainty: $\sim 2.5\%$.
- ▶ Increasing the flux uncertainty is a game that one can play, but there are only guesses, e.g. $\sim 5\%$ [Hayes and Vogel, ARNPS 66 (2016) 219].
- ▶ Bottom line: the status of the reactor anomaly is controversial.

NEOS

[PRL 118 (2017) 121802 (arXiv:1610.05134)]





- ▶ Hanbit Nuclear Power Complex in Yeong-gwang, Korea.
- ▶ Thermal power of 2.8 GW.
- ▶ Detector: a ton of Gd-loaded liquid scintillator in a gallery approximately 24 m from the reactor core.
- ▶ The measured antineutrino event rate is 1976 per day with a signal to background ratio of about 22.

DANSS


[Danilov @ Solvay Workshop, 1 December 2017, and La Thuile 2018, 3 March 2018]

Detector of reactor AntiNeutrino based on Solid Scintillator




- ▶ Installed on a movable platform under a 3 GW reactor.
- ▶ Large neutrino flux.
- ▶ Reactor shielding of cosmic rays.
- ▶ Variable source-detector distance with the same detector!

$$\begin{aligned} \text{Down} &= 12.7 \text{ m} \\ \text{Up} &= 10.7 \text{ m} \end{aligned}$$



Model-Independent $\bar{\nu}_e$ SBL Oscillations

[Gariazzo, CG, Laveder, Li, PLB 782 (2018) 13, arXiv:1801.06467]



$\sim 3.7\sigma$

$$\Delta m_{41}^2 = 1.29 \pm 0.03$$


$$\sin^2 2\vartheta_{ee} = 0.049 \pm 0.011$$

$$\sin^2 \vartheta_{14} = |U_{e4}|^2$$

$$\sin^2 \vartheta_{14} = 0.012 \pm 0.003$$


$$\sin^2 \vartheta_{13} = 0.022 \pm 0.001$$

Comparison with the Reactor and Gallium Anomalies




- ▶ 3 σ agreement.
- ▶ 2 σ tension.
- ▶ Small overestimate of the reactor fluxes.
- ▶ Small overestimate of the GALLEX and SAGE efficiencies.

Global Model-Independent ν_e and $\bar{\nu}_e$ Disappearance



- ▶ NEOS and DANSS.
- ▶ Reactor rates with free ²³⁵U and ²³⁹Pu fluxes: r_{235} and r_{239} .
- ▶ Gallium data with free GALLEX and SAGE efficiencies: η_G and η_S .
- ▶ New reactor experiments: STEREO, Neutrino-4, SoLiD, PROSPECT
- ▶ Kinematical ν_4 mass measurement: KATRIN

[See also Dentler, Hernandez-Cabezudo, Kopp, Machado, Maltoni, Martinez-Soler, Schwetz, arXiv:1803.10661]




- Indication of $r_{235} < 1$.
- Likely small overestimate of the GALLEX and SAGE efficiencies.

LSND

[PRL 75 (1995) 2650; PRC 54 (1996) 2685; PRL 77 (1996) 3082; PRD 64 (2001) 112007]


$$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$$

$$20 \text{ MeV} \leq E \leq 52.8 \text{ MeV}$$



$$\Delta m_{SBL}^2 \gtrsim 0.1 \text{ eV}^2 \gg \Delta m_{ATM}^2$$

- Well-known and pure source of $\bar{\nu}_\mu$



Well-known detection process of $\bar{\nu}_e$

- $\approx 3.8\sigma$ excess
- But signal not seen by KARMEN at $L \simeq 18 \text{ m}$ with the same method

[PRD 65 (2002) 112001]

MiniBooNE

$L \simeq 541 \text{ m}$



$200 \text{ MeV} \leq E \lesssim 3 \text{ GeV}$

$$\nu_\mu \rightarrow \nu_e$$

[PRL 102 (2009) 101802]


$$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$$

[PRL 110 (2013) 161801]




- ▶ Purpose: check LSND signal.
- ▶ LSND signal: $E > 475 \text{ MeV}$.
- ▶ Different L and E .
- ▶ Agreement with LSND signal?
- ▶ Similar L/E (oscillations).
- ▶ Low-energy anomaly \Rightarrow MicroBooNE
- ▶ No money, no Near Detector.
- ▶ Pragmatic Approach: $E > 475 \text{ MeV}$.

$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ and $\nu_\mu \rightarrow \nu_e$ Appearance



ν_μ and $\bar{\nu}_\mu$ Disappearance




3+1 Appearance-Disappearance Tension

$$\nu_e \text{ DIS} \\ \sin^2 2\vartheta_{ee} \simeq 4|U_{e4}|^2$$

$$\nu_\mu \text{ DIS} \\ \sin^2 2\vartheta_{\mu\mu} \simeq 4|U_{\mu 4}|^2$$

$$\nu_\mu \rightarrow \nu_e \text{ APP} \\ \sin^2 2\vartheta_{e\mu} = 4|U_{e4}|^2|U_{\mu 4}|^2 \simeq \frac{1}{4} \sin^2 2\vartheta_{ee} \sin^2 2\vartheta_{\mu\mu}$$





- ▶ $\nu_\mu \rightarrow \nu_e$ is quadratically suppressed!
- ▶ Global Fit without MINOS+
 $\chi^2_{\text{PG}}/\text{NDF}_{\text{PG}} = 5.5/2 \Rightarrow \text{GoF}_{\text{PG}} = 6.5\%$
- ▶ Similar tension in 3+2, 3+3, ..., 3+N_s



[CG, Zavaini, MPLA 31 (2015) 1650003]

New Bound from MINOS+

[arXiv:1710.06488]



Effects of MINOS+



- $\chi^2_{\text{PG}}/\text{NDF}_{\text{PG}} = 14.6/2 \Rightarrow \text{GoF}_{\text{PG}} = 0.07\%$ ← Intolerable tension!
- The MINOS+ bound (if correct) disfavors the LSND $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ signal.

[See also Dentler, Hernandez-Cabezudo, Kopp, Machado, Maltoni, Martinez-Soler, Schwetz, arXiv:1803.10661]

Conclusions

- ▶ Exciting model-independent indication of light sterile neutrinos at the eV scale from the NEOS and DANSS experiments \Rightarrow New Physics beyond the Standard Model!
- ▶ Agreement with the Reactor and Gallium Anomalies \Rightarrow Needed revision of the ^{235}U calculation and small decrease of the GALLEX and SAGE efficiencies.
- ▶ Can be checked in the near future by the reactor experiments STEREO, Neutrino-4, SoLid, PROSPECT.
- ▶ Independent tests through effect of m_4 in β -decay (**KATRIN**) and $\beta\beta_{0\nu}$ -decay.
- ▶ The MINOS+ bound (if correct) disfavors the LSND $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ signal.